

**EXTENDED RANGE FORECAST OF ATLANTIC SEASONAL HURRICANE
ACTIVITY AND LANDFALL STRIKE PROBABILITY FOR 2009**

We foresee slightly below-average activity for the 2009 Atlantic hurricane season. We have reduced our seasonal forecast from our early April prediction. We anticipate a slightly below-average probability of United States and Caribbean major hurricane landfall.

(as of 2 June 2009)

By Philip J. Klotzbach¹ and William M. Gray²

This forecast as well as past forecasts and verifications are available via the World Wide Web at <http://hurricane.atmos.colostate.edu/Forecasts>

Emily Wilmsen, Colorado State University Media Representative, (970-491-6432) is available to answer various questions about this forecast

Department of Atmospheric Science
Colorado State University
Fort Collins, CO 80523
Email: amie@atmos.colostate.edu

¹ Research Scientist

² Professor Emeritus of Atmospheric Science

Why issue extended-range forecasts for seasonal hurricane activity?

We are frequently asked this question. Our answer is that it is possible to say something about the probability of the coming year's hurricane activity which is superior to climatology. The Atlantic basin has the largest year-to-year variability of any of the global tropical cyclone basins. People are curious to know how active the upcoming season is likely to be, particularly if you can show hindcast skill improvement over climatology for many past years.

Everyone should realize that it is impossible to precisely predict this season's hurricane activity in early June. There is, however, much curiosity as to how global ocean and atmosphere features are presently arranged as it regards to the probability of an active or inactive hurricane season for the coming year. Our early June statistical forecast methodology shows strong evidence over 58 past years that significant improvement over climatology can be attained. The model correctly predicted an above-average season in 2008. We would never issue a seasonal hurricane forecast unless we had a statistical model developed over a long hindcast period which showed significant skill over climatology.

We issue these forecasts to satisfy the curiosity of the general public and to bring attention to the hurricane problem. There is a general interest in knowing what the odds are for an active or an inactive season. One must remember that our forecasts are based on the premise that those global oceanic and atmospheric conditions which preceded comparatively active or inactive hurricane seasons in the past provide meaningful information about similar trends in future seasons. This is not always true for individual seasons. It is also important that the reader appreciate that these seasonal forecasts are based on statistical schemes which, owing to their intrinsically probabilistic nature, will fail in some years. Moreover, these forecasts do not specifically predict where within the Atlantic basin these storms will strike. The probability of landfall for any one location along the coast is very low and reflects the fact that, in any one season, most U.S. coastal areas will not feel the effects of a hurricane no matter how active the individual season is. However, all coastal residents should prepare for hurricane season every year, since landfalling tropical cyclones can devastate communities in inactive or active seasons. It only takes one landfalling system to make this a very active season for you.

ATLANTIC BASIN SEASONAL HURRICANE FORECAST FOR 2009

Forecast Parameter and 1950-2000 Climatology (in parentheses)	Issue Date 10 December 2008	Issue Date 9 April 2009	Issue Date 2 June 2009
Named Storms (NS) (9.6)	14	12	11
Named Storm Days (NSD) (49.1)	70	55	50
Hurricanes (H) (5.9)	7	6	5
Hurricane Days (HD) (24.5)	30	25	20
Major Hurricanes (MH) (2.3)	3	2	2
Major Hurricane Days (MHD) (5.0)	7	5	4
Accumulated Cyclone Energy (ACE) (96.1)	125	100	85
Net Tropical Cyclone Activity (NTC) (100%)	135	105	90

PROBABILITIES FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE
LANDFALL ON EACH OF THE FOLLOWING UNITED STATES COASTAL
AREAS:

- 1) Entire U.S. coastline - 48% (average for last century is 52%)
- 2) U.S. East Coast Including Peninsula Florida - 28% (average for last century is 31%)
- 3) Gulf Coast from the Florida Panhandle westward to Brownsville - 28% (average for last century is 30%)

PROBABILITY FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE
TRACKING INTO THE CARIBBEAN (10-20°N, 60-88°W)

- 1) 39% (average for last century is 42%)

ABSTRACT

Information obtained through May 2009 indicates that the 2009 Atlantic hurricane season will be slightly less active than the average 1950-2000 season. We estimate that 2009 will have about 5 hurricanes (average is 5.9), 11 named storms (average is 9.6), 50 named storm days (average is 49.1), 20 hurricane days (average is 24.5), 2 major (Category 3-4-5) hurricanes (average is 2.3) and 4 major hurricane days (average is 5.0). The probability of U.S. major hurricane landfall and Caribbean major hurricane activity is estimated to be slightly below the long-period average. We expect Atlantic basin Net Tropical Cyclone (NTC) activity in 2009 to be approximately 90 percent of the long-term average. We have decreased our seasonal forecast from early April.

This forecast is based on an extended-range early June statistical prediction scheme that utilizes 58 years of past data. Analog predictors are also utilized. The influence of El Niño conditions is implicit in these predictor fields, and therefore we do not utilize a specific ENSO forecast as a predictor.

We expect current neutral ENSO conditions to persist or perhaps transition to weak El Niño conditions by the most active portion of this year's hurricane season (August-October). If El Niño conditions develop, it would tend to increase the levels of vertical wind shear and decrease the levels of Atlantic hurricane activity. Another reason for our forecast reduction is due to the persistence of anomalously cool sea surface temperatures in the tropical Atlantic. Cooler waters are associated with dynamic and thermodynamic factors that are less conducive for an active Atlantic hurricane season. Another factor in our forecast reduction is the stronger-than-normal Azores High during April-May. Stronger high pressure typically results in stronger trade winds that are commonly associated with less active hurricane seasons.

Although we have been in an active multi-decadal Atlantic Basin hurricane era since 1995, it is not unusual to have a few below-average years within an active multi-decadal period. Likewise, it is not unusual to have a few above-average years within an inactive multi-decadal period. We expect the active Atlantic hurricane era that we have been in since 1995 to continue for the next 10-15 years.

Notice of Author Changes

By William Gray

The order of the authorship of these forecasts was reversed in 2006 from Gray and Klotzbach to Klotzbach and Gray. After 22 years (1984-2005) of making these forecasts, it was appropriate that I step back and have Phil Klotzbach assume the primary responsibility for our project's seasonal, monthly and landfall probability forecasts. Phil has been a member of my research project for the last nine years and was second author on these forecasts from 2001-2005. I have greatly profited and enjoyed our close personal and working relationships.

Phil is now devoting much more time to the improvement of these forecasts than I am. I am now giving more of my efforts to the global warming issue and in synthesizing my projects' many years of hurricane and typhoon studies.

Phil Klotzbach is an outstanding young scientist with a superb academic record. I have been amazed at how far he has come in his knowledge of hurricane prediction since joining my project in 2000. I foresee an outstanding future for him in the hurricane field. He is currently making many new seasonal and monthly forecast innovations that are improving our forecasts. The success of last year's seasonal forecasts is an example. Phil was awarded his Ph.D. degree in 2007. He is currently spending most of his time working towards better understanding and improving these Atlantic basin hurricane forecasts.

Acknowledgment

We are grateful to the National Science Foundation (NSF) for providing partial support for the research necessary to make these forecasts. We also thank the GeoGraphics Laboratory at Bridgewater State College (MA) for their assistance in developing the United States Landfalling Hurricane Probability Webpage (available online at <http://www.e-transit.org/hurricane>).

The second author gratefully acknowledges the valuable input to his CSU seasonal forecast research project over many years by former project members and now colleagues Chris Landsea, John Knaff and Eric Blake. We also thank Professors Paul Mielke and Ken Berry of Colorado State University for much statistical analysis and advice over many years. We also thank Bill Thorson for technical advice and assistance.

DEFINITIONS

Accumulated Cyclone Energy – (ACE) A measure of a named storm's potential for wind and storm surge destruction defined as the sum of the square of a named storm's maximum wind speed (in 10^4 knots²) for each 6-hour period of its existence. The 1950-2000 average value of this parameter is 96.

Atlantic Basin – The area including the entire North Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico.

El Niño – (EN) A 12-18 month period during which anomalously warm sea surface temperatures occur in the eastern half of the equatorial Pacific. Moderate or strong El Niño events occur irregularly, about once every 3-7 years on average.

Hurricane – (H) A tropical cyclone with sustained low-level winds of 74 miles per hour (33 ms^{-1} or 64 knots) or greater.

Hurricane Day – (HD) A measure of hurricane activity, one unit of which occurs as four 6-hour periods during which a tropical cyclone is observed or estimated to have hurricane intensity winds.

Main Development Region (MDR) – An area in the tropical Atlantic where a majority of major hurricanes form, defined as 10-20°N, 70-20°W.

Major Hurricane – (MH) A hurricane which reaches a sustained low-level wind of at least 111 mph (96 knots or 50 ms^{-1}) at some point in its lifetime. This constitutes a category 3 or higher on the Saffir/Simpson scale.

Major Hurricane Day – (MHD) Four 6-hour periods during which a hurricane has an intensity of Saffir/Simpson category 3 or higher.

Named Storm – (NS) A hurricane, a tropical storm or a sub-tropical storm.

Named Storm Day – (NSD) As in HD but for four 6-hour periods during which a tropical or sub-tropical cyclone is observed (or is estimated) to have attained tropical storm intensity winds.

NTC – Net Tropical Cyclone Activity – Average seasonal percentage mean of NS, NSD, H, HD, IH, IHD. Gives overall indication of Atlantic basin seasonal hurricane activity. The 1950-2000 average value of this parameter is 100.

QBO – Quasi-Biennial Oscillation – A stratospheric (16 to 35 km altitude) oscillation of equatorial east-west winds which vary with a period of about 26 to 30 months or roughly 2 years; typically blowing for 12-16 months from the east, then reversing and blowing 12-16 months from the west, then back to easterly again.

Saffir/Simpson (S-S) Category – A measurement scale ranging from 1 to 5 of hurricane wind and ocean surge intensity. One is a weak hurricane; whereas, five is the most intense hurricane.

SOI – Southern Oscillation Index – A normalized measure of the surface pressure difference between Tahiti and Darwin.

SST(s) – Sea Surface Temperature(s)

SSTA(s) – Sea Surface Temperature(s) Anomalies

Tropical Cyclone – (TC) A large-scale circular flow occurring within the tropics and subtropics which has its strongest winds at low levels; including hurricanes, tropical storms and other weaker rotating vortices.

Tropical North Atlantic (TNA) index – A measure of sea surface temperatures in the area from 5.5-23.5°N, 57.5-15°W.

Tropical Storm – (TS) A tropical cyclone with maximum sustained winds between 39 (18 ms^{-1} or 34 knots) and 73 (32 ms^{-1} or 63 knots) miles per hour.

ZWA – Zonal Wind Anomaly – A measure of the upper level (~200 mb) west to east wind strength. Positive anomaly values mean winds are stronger from the west or weaker from the east than normal.

1 knot = 1.15 miles per hour = 0.515 meters per second

1 Introduction

This is the 26th year in which the CSU Tropical Meteorology Project has made forecasts of the upcoming season's Atlantic basin hurricane activity. Our research team has shown that a sizable portion of the year-to-year variability of Atlantic tropical cyclone (TC) activity can be hindcast with skill exceeding climatology. These forecasts are based on a statistical methodology derived from 58 years of past data. Qualitative adjustments are added to accommodate additional processes which may not be explicitly represented by our statistical analyses. These evolving forecast techniques are based on a variety of climate-related global and regional predictors previously shown to be related to the forthcoming seasonal Atlantic basin tropical cyclone activity and landfall probability. We believe that seasonal forecasts must be based on methods that show significant hindcast skill in application to long periods of prior data. It is only through hindcast skill that one can demonstrate that seasonal forecast skill is possible. This is a valid methodology provided that the atmosphere continues to behave in the future as it has in the past.

The best predictors do not necessarily have the best individual correlations with hurricane activity. The best forecast parameters are those that explain the portion of the variance of seasonal hurricane activity that is not associated with the other forecast variables. It is possible for an important hurricane forecast parameter to show little direct relationship to a predictand by itself but to have an important influence when included with a set of 2-3 other predictors.

A direct correlation of a forecast parameter may not be the best measure of the importance of this predictor to the skill of a 2-3 parameter forecast model. This is the nature of the seasonal or climate forecast problem where one is dealing with a very complicated atmosphere-ocean system that is highly non-linear. There is a maze of changing physical linkages between the many variables. These linkages can undergo unknown changes from weekly to decadal time scales. It is impossible to understand how all these processes interact with each other. No one can completely understand the full complexity of the atmosphere-ocean system. But, it is still possible to develop a reliable statistical forecast scheme which incorporates a number of the climate system's non-linear interactions. Any seasonal or climate forecast scheme must show significant hindcast skill before it is used in real-time forecasts.

2 June Forecast Methodology

We developed a new June forecast scheme which was used for the first time last year. This scheme worked out quite well in predicting a very active season last year. Complete details on the earlier June forecast schemes used from 1995-2007 are available in our June 2008 forecast ([Klotzbach and Gray 2008](#)).

2.1 Current June Statistical Forecast Scheme

We have found that using two spring predictors and our early April hindcast, we can obtain early June hindcasts that show considerable skill over the period from 1950-2007. This new forecast model also provided a very accurate prediction for the 2008 hurricane season.

This new scheme was created by evaluating the two spring predictors using least-squared regression. The resulting hindcasts were then ranked in order from 1 (the highest value) to 58 (the lowest value). Then the resulting preliminary June NTC hindcast rank was adjusted to the final June NTC hindcast by using the following method. We ranked the April NTC hindcasts in a similar manner as was done with early June (i.e., from 1 to 58). Then the final June NTC hindcast rank was derived by computing the following equation:

$$\text{Final June NTC Hindcast Rank} = 0.5 * (\text{Preliminary June NTC Hindcast Rank}) + 0.5 * (\text{Final April NTC Hindcast Rank}).$$

The final NTC hindcast was obtained by taking the final June NTC hindcast rank and assigning the observed NTC value for that rank. For example, if the final June NTC hindcast rank was 10 (the 10th highest rank), the NTC value assigned for the prediction would be the 10th highest observed rank, which in this case would be 166 NTC units. Since there is considerable uncertainty at this extended lead time as to final forecast values, final hindcast values are constrained to be between 40 and 200 NTC units.

Using the ranking method to arrive at our final forecast values is a new statistical forecasting approach for us. We find that using this method improves the hindcast skill of our forecasts somewhat (approximately 4-10%) and also allows for improved predictability of outliers. For example, simply by ranking our December hindcasts and assigning observed NTC values to those ranks improves our hindcast skill (as measured by variance explained) in early December from 45% to 49%.

As mentioned before, our new statistical scheme shows enhanced levels of hindcast skill, explaining 66 percent of the variance from 1950-2007 and 79 percent of the variance from 1995-2007. We believe that we have solid physical links between these predictors and the upcoming Atlantic basin hurricane season.

Table 1 displays our early June hindcasts for 1950-2007 using the new statistical scheme, while Figure 1 displays observations versus NTC hindcasts. Our early June hindcasts have correctly predicted above- or below-average seasons in 46 out of 58 hindcast years (79%). These hindcasts have had a smaller error than climatology in 36 out of 58 years (62%). Our average hindcast error is 27 NTC units, compared with 44 NTC units for climatology. This scheme also shows considerable stability when broken in half, explaining 56 percent of the variance from 1950-1978 and 77 percent of the variance from 1979-2007. The scheme has shown remarkable skill over the past 28 years (since 1980). The model has had a smaller error than climatology in 20 out of the last 28 years (71%). This new scheme is also well-tuned to the multi-decadal active hurricane periods from 1950-1969 and 1995-2007 versus the inactive hurricane period from 1970-

1994 (Table 2). Figure 2 displays the locations of the two 1 June (using April-May data) predictors used in this scheme in map form. Please refer to Figure 2 of our early April forecast for locations of predictors used in our early April prediction scheme. Table 3 lists the three (two new spring predictors and our early April prediction) predictors that are utilized for this year's June forecast. A more extensive discussion of current conditions in the Pacific and Atlantic basins is provided in Sections 5 and 6, respectively.

Table 1: Observed versus hindcast NTC for 1950-2007 using the current June scheme. Average errors for hindcast NTC and climatological NTC predictions are given without respect to sign. Red bold-faced years in the “Hindcast NTC” column (2) are years that we did not go the right way, while red bold-faced years in the “Hindcast improvement over Climatology” column (5) are years that we did not beat climatology. The hindcast went the right way with regards to an above- or below-average season in 46 out of 58 years (79%), while hindcast improvement over climatology occurred in 36 out of 58 years (62%). The real-time forecast for 2008 is also listed.

Year	(1) Observed NTC	(2) Hindcast NTC	(3) Observed minus Hindcast	(4) Observed minus Climatology	(5) Hindcast improvement over Climatology
1950	230	200	30	130	100
1951	115	82	34	15	-18
1952	93	173	-80	-7	-73
1953	116	200	-84	16	-69
1954	124	130	-6	24	18
1955	188	188	0	88	88
1956	66	98	-32	-34	2
1957	82	93	-12	-18	7
1958	133	134	-1	33	32
1959	94	109	-15	-6	-9
1960	92	129	-37	-8	-29
1961	211	200	11	111	100
1962	32	97	-65	-68	3
1963	111	82	29	11	-18
1964	160	133	27	60	33
1965	82	124	-42	-18	-24
1966	134	116	19	34	16
1967	93	82	11	-7	-5
1968	39	52	-13	-61	48
1969	150	106	44	50	6
1970	62	72	-11	-38	28
1971	91	62	29	-9	-21
1972	27	40	-13	-73	60
1973	50	92	-42	-50	8
1974	72	45	28	-28	0
1975	89	66	23	-11	-12
1976	82	51	31	-18	-12
1977	45	40	5	-55	51
1978	83	40	43	-17	-26
1979	92	40	52	-8	-43
1980	129	80	49	29	-20
1981	109	85	24	9	-15
1982	35	50	-14	-65	50
1983	31	40	-9	-69	60
1984	74	115	-41	-26	-15
1985	106	92	14	6	-8
1986	37	40	-3	-63	60
1987	46	91	-46	-54	9
1988	118	93	25	18	-7
1989	130	160	-30	30	0
1990	98	94	4	-2	-2
1991	57	46	11	-43	32
1992	64	40	24	-36	12
1993	52	83	-31	-48	17
1994	35	57	-22	-65	43
1995	222	185	37	122	85
1996	192	192	0	92	92
1997	51	64	-13	-49	36
1998	166	134	31	66	34
1999	185	200	-15	85	70
2000	134	150	-16	34	18
2001	129	111	18	29	11
2002	80	74	6	-20	14
2003	173	129	44	73	29
2004	228	166	63	128	66
2005	273	200	73	173	100
2006	85	118	-33	-15	-18
2007	99	89	10	-1	-9
Average	106	104	27	44	+17
2008	162	160	2	62	+60

Hindcast vs. Observed NTC - 1 June - Rank Prediction Method




Figure 1: Observed versus hindcast values of NTC for 1950-2007.

Table 2: Hindcast versus observed average NTC for active vs. inactive multi-decadal periods.

<i>Years</i>	<i>Average Hindcast NTC</i>	<i>Average Observed NTC</i>
1950-1969 (Active)	126	117
1970-1994 (Inactive)	69	72
1995-2007 (Active)	139	155

New June Forecast Predictors




Figure 2: Location of spring predictors for our June extended-range statistical prediction for the 2009 hurricane season.

Table 3: Listing of 1 June 2009 predictors using the June statistical model for the 2009 hurricane season. A plus (+) means that positive values of the parameter indicate increased hurricane activity during the following year.

Predictor	2009 Forecast Values
1) Subtropical Atlantic Index (+): April-May SST (20-50°N, 15-30°W) (+) & May SLP (10-35°N, 10-40°W) (-)	-0.4 SD
2) April-May 200 MB U (5-25°S, 50-90°E) (-)	+0.2 SD
3) Early April Hindcast (+)	110 NTC

There is also extended-range forecast skill from 1 June for United States hurricane landfall probabilities. In the 15 out of 58 years where our current hindcast scheme forecast NTC values that were above 130, we had more than twice as many hurricane (41 versus 20) and more than three times as many major hurricane (17 versus 5) landfalls along the U.S. coastline when compared with the 15 out of 58 years where our hindcast scheme gave NTC values that were below 65. For the Florida Peninsula and the U.S. East Coast, the ratio between NTC hindcast values greater than 130 and below 65 are 25 to 9 for hurricanes and 9 to 1 for major hurricanes – a remarkable difference.

2.2 Physical Associations among Predictors Listed in Table 3

The locations and brief descriptions of the two spring predictors for our early June statistical forecast are now discussed. It should be noted that both forecast parameters correlate significantly with seasonal physical features that are known to be favorable for elevated levels of hurricane activity. These factors are primarily related to August-October vertical wind shear in the Atlantic Main Development Region (MDR) from 10-20°N, 20-70°W as shown in Figure 3.




Figure 3: Vertical wind profile typically associated with (a) inactive Atlantic basin hurricane seasons and (b) active Atlantic basin hurricane seasons. Note that (b) has reduced levels of vertical wind shear.

For each of these predictors, we display a four-panel figure showing linear correlations between values of each predictor and August-October values of sea surface temperature, sea level pressure, 200 mb zonal wind, and 925 mb zonal wind, respectively. In general, higher values of SSTA, lower values of SLPA, anomalous westerlies at 925 mb and anomalous easterlies at 200 mb are associated with active Atlantic basin hurricane seasons.

For more information about the predictors utilized in our early April statistical forecast (used as 50% of our early June forecast), please refer to our early April 2009 forecast:

1. Subtropical Atlantic Index (+): April-May SST (20-50°N, 15-30°W) (+) & May SLP (10-35°N, 10-40°W) (-)

A combination of above-normal sea surface temperatures (SSTs) in the eastern subtropical Atlantic and lower-than-normal sea level pressures in the subtropical Atlantic is associated with a weakened Azores high and reduced trade wind strength during the late spring (Knaff 1997). This combined index in April-May is strongly correlated with weaker trade winds, lower-than-normal sea level pressures and above-normal SSTs in the tropical Atlantic during the following August-October period (Figure 4). All three of these August-October features are commonly associated with active Atlantic basin hurricane seasons, through reductions in vertical wind shear, increased vertical instability and increased surface latent and sensible heat fluxes, respectively. Stronger-than-normal values of this index correlate quite well (~0.5) with active Atlantic basin tropical cyclone seasons.

Predictor 2. April-May 200 MB U in the South Indian Ocean (-)

(5-25°S, 50-90°E)

Upper-level easterly wind anomalies in the South Indian Ocean during April-May are associated with anomalously low sea level pressure and anomalous upper-level divergence in the western tropical Pacific and anomalously high sea level pressure and anomalous upper-level convergence in the eastern tropical Pacific. These features are associated with an active Walker Circulation, which is typically observed in cool ENSO years. Figure 5 displays the significant correlations that are achieved between values of this predictor in April-May and August-October sea surface temperatures, sea level pressure and 925 and 200 mb zonal wind anomalies, respectively. Note the anomalous easterly winds that are typically observed at upper levels over the tropical Atlantic and Caribbean in August-October when upper-level easterlies exist in the South Indian Ocean in April-May. These anomalous easterlies, combined with anomalous westerlies at 925 mb, reduce vertical wind shear across the tropical Atlantic providing a more favorable environment for tropical cyclone formation and intensification. Predictor values have been trending slightly more positive in this region since the 1950s. We have removed the trend in zonal wind anomalies from our predictor calculations to avoid a potentially non-physical lowering of forecast values, as there is some uncertainty as to the quality of the NCEP/NCAR reanalysis data for upper-level winds in the 1950s.

This predictor is located in an area that has not been considered in our previous early June forecasts. Since this predictor is new, we have gone through extensive testing to make sure that the predictor is valid. The predictor shows considerable stability when evaluated over both the 1950-1989 period and the 1990-2007 period. It correlates with NTC at -0.57 over the period from 1950-1989 and correlates with NTC at -0.60 over the period from 1990-2007. The correlation with NTC over the full time period from 1950-2007 is -0.59 (Figure 6).

When we examined the top 10 years when the predictor had its highest values and compared them with the bottom 10 years when the predictor had its lowest values, considerable differences were evident. In the 10 years when the zonal winds had their largest easterly anomalies, an average of 140 NTC units were observed, compared with an average of only 60 NTC units in the 10 years when the zonal winds had their largest westerly anomalies (Figure 7).




Figure 4: Linear correlations between the subtropical Atlantic index (Predictor 1) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 925 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations are known to be favorable for enhanced hurricane activity.




Figure 5: Linear correlations between April-May 200 mb U in the South Indian Ocean (Predictor 2) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 925 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations are known to be favorable for enhanced hurricane activity. Zonal wind values have been multiplied by -1 to allow for easy comparison with Figure 4.




Figure 6: April-May values of Predictor 2 (pink line) and Atlantic basin NTC activity (blue line). Note the strong negative correlation between the two curves.




Figure 7: Top 10/bottom 10 ratios for Predictor 2. Note that much larger values of NTC were observed when Predictor 2 had anomalous easterly winds.

3 Forecast Uncertainty

One of the questions that we are asked fairly frequently regarding our seasonal hurricane predictions is the degree of uncertainty that is involved. Obviously, our predictions are our best estimate, but there is with all forecasts an uncertainty as to how well they will verify.

Table 4 provides our early June forecasts, with error bars (based on one standard deviation of absolute errors) as calculated from hindcasts over the 1990-2007 period, using equations developed over the 1950-1989 period. We typically expect to see 2/3 of our forecasts verify within one standard deviation of observed values, with 95% of forecasts verifying within two standard deviations of observed values.

Table 4: Model hindcast error and our 2009 hurricane forecast. Uncertainty ranges are given in one standard deviation (SD) increments.

Parameter	Hindcast Error (SD)	2009 Forecast	Uncertainty Range – 1 SD (67% of Forecasts Likely in this Range)
Named Storms (NS)	3.8	11	7.2 – 14.8
Named Storm Days (NSD)	18.3	50	31.7 – 68.3
Hurricanes (H)	2.1	5	3.9 – 8.1
Hurricane Days (HD)	9.0	20	11.0 – 29.0
Major Hurricanes (MH)	1.2	2	0.8 – 3.2
Major Hurricane Days (MHD)	4.5	4	0.0 – 8.5
Accumulated Cyclone Energy (ACE)	39	85	46 – 124
Net Tropical Cyclone (NTC) Activity	37	90	53 – 127

4 Analog-Based Predictors for 2009 Hurricane Activity

Certain years in the historical record have global oceanic and atmospheric trends which are substantially similar to 2009. These years also provide useful clues as to likely trends in activity that the forthcoming 2009 hurricane season may bring. For this early June extended range forecast, we determine which of the prior years in our database have distinct trends in key environmental conditions which are similar to current April-May 2009 conditions. Table 5 lists our analog selections.

We select prior hurricane seasons since 1949 which have similar atmospheric-oceanic conditions to those currently being experienced. We searched for years that were generally characterized by neutral to slightly warm ENSO conditions, slightly below-average tropical Atlantic SSTs and above-average far North Atlantic SSTs during April-May.

There were five hurricane seasons since 1949 with characteristics most similar to what we observed in April-May 2009. The best analog years that we could find for the 2009 hurricane season were 1959, 1960, 1965, 2001 and 2002. We anticipate that 2009 seasonal hurricane activity will have activity in line with what was experienced in the average of these five years. We believe that 2009 will have slightly below-average activity in the Atlantic basin.

Table 5: Best analog years for 2009 with the associated hurricane activity listed for each year.

Year	NS	NSD	H	HD	MH	MHD	ACE	NTC
1959	11	40.00	7	22.00	2	4.25	78	96
1960	7	29.50	4	18.25	2	9.75	88	93
1965	6	39.50	4	27.25	1	7.50	84	86
2001	15	68.75	9	25.50	4	4.25	110	135
2002	12	57.00	4	10.75	2	3.00	67	83
Mean	10.2	47.0	5.6	20.8	2.2	5.8	86	99
2009 Forecast	11	50	5	20	2	4	85	90

5 ENSO

Neutral ENSO conditions have occurred so far this spring. SSTs have generally warmed over the past two months. They are now slightly above average across the eastern and central tropical Pacific. Table 6 displays March and May SST anomalies for several Nino regions. Note that all four regions have experienced warming since March, with more warming occurring in the eastern Pacific.

Table 6: March and May SST anomalies for Nino 1+2, Nino 3, Nino 3.4, and Nino 4, respectively. May-March SST anomaly differences are also provided.

Region	March SST Anomaly (°C)	May SST Anomaly (°C)	March – January SST Anomaly (°C)
Nino 1+2	-0.1	0.5	+0.6
Nino 3	-0.6	0.4	+1.0
Nino 3.4	-0.2	0.4	+0.6
Nino 4	0.0	0.4	+0.4

As was the situation last year, the big question is whether this current observed warming will continue through this year’s hurricane season. Several of the dynamical ENSO models are calling for a weak or moderate ENSO event during the peak of this year’s hurricane season (August-October), while most statistical models are calling for the continuation of neutral conditions (Figure 8).

Model Forecasts of ENSO from *May 2009*




Figure 8: ENSO forecasts from various statistical and dynamical models. Figure courtesy of the International Research Institute (IRI). Most of the dynamical models are now calling for an El Niño event by August-October, while most of the statistical models are calling for the persistence of neutral conditions.

We believe that there is a slightly greater chance of a weak El Niño developing this summer/fall than there was in early April. SST anomalies have continued to moderate since then, and the current wind/ocean heat content pattern could likely sustain a weak El Niño. At this point, we believe there is an approximately 70% chance of a weak El Niño (ASO Nino 3.4 > 0.5°C) by the August-October period. This increased potential for a weak El Niño is one of the reasons that we have reduced our forecast from early April. El Niños typically increase levels of vertical wind shear in the tropical Atlantic, causing detrimental conditions for Atlantic tropical cyclone formation and intensification.

6 Current Atlantic Basin Conditions

Conditions in the Atlantic have continued to remain less favorable for an active season. Figure 9 displays the SST anomaly difference between May 2009 and November 2008. Note the strong anomalous cooling that has occurred across the Main Development Region. Current Tropical North Atlantic index (defined as 5.5-23.5°N, 57.5-15°W) SST anomaly values of approximately -0.4°C are the lowest that have been observed since June-July 1994. High sea level pressure anomalies and stronger-than-normal trade winds in the eastern subtropical Atlantic have been observed for the past couple of months, indicating that these cool SST anomalies will likely persist due to enhanced evaporation and upwelling. Cooler-than-normal waters are detrimental for tropical cyclone formation and intensification because they provide less latent and sensible heat flux. In addition, an anomalously cool tropical Atlantic is usually associated with higher sea level pressure values and stronger-than-normal trade winds, indicating a more stable atmosphere with increased levels of vertical wind shear. Cooler SSTs and a stronger-than-normal April-May Azores High are also indicative of stronger-than-normal Atlantic vertical wind shear.




Figure 9: May 2009 – November 2008 SST anomaly difference across the Atlantic. In general, the Atlantic has cooled considerably over the past six months.

7 Adjusted 2009 Forecast

Table 7 shows our final adjusted early June forecast for the 2009 season which is a combination of our statistical scheme, our analog forecast and qualitative adjustments for other factors not explicitly contained in any of these schemes. Our statistical forecast and our analog forecast indicate activity at near-average levels. We foresee a slightly below-average Atlantic basin hurricane season due to currently-observed unfavorable conditions in the tropical Atlantic. Another reason for our reduction in our early June forecast from our forecast of early April is our belief in an increased probability of a weak El Niño developing.

Table 7: Summary of our early June statistical forecast, our analog forecast and our adjusted final forecast for the 2009 hurricane season.

Forecast Parameter and 1950-2000 Climatology (in parentheses)	Statistical Scheme	Analog Scheme	Adjusted Final Forecast
Named Storms (9.6)	10.0	10.2	11
Named Storm Days (49.1)	49.1	47.0	50
Hurricanes (5.9)	5.9	5.6	5
Hurricane Days (24.5)	23.2	20.8	20
Intense Hurricanes (2.3)	2.5	2.2	2
Intense Hurricane Days (5.0)	5.7	5.8	4
Accumulated Cyclone Energy Index (96.1)	94	86	85
Net Tropical Cyclone Activity (100%)	103	99	90

8 Landfall Probabilities for 2009

A significant focus of our recent research involves efforts to develop forecasts of the probability of hurricane landfall along the U.S. coastline. Whereas individual hurricane landfall events cannot be accurately forecast months in advance, the total seasonal probability of landfall can be forecast with statistical skill. With the observation that, statistically, landfall is a function of varying climate conditions, a probability specification has been developed through statistical analyses of all U.S. hurricane and named storm landfall events during the 20th century (1900-1999). Specific landfall probabilities can be given for all tropical cyclone intensity classes for a set of distinct U.S. coastal regions.

Net landfall probability is shown linked to the overall Atlantic basin Net Tropical Cyclone activity (NTC; see Table 8). NTC is a combined measure of the year-to-year mean of six indices of hurricane activity, each expressed as a percentage difference from the long-term average. Long-term statistics show that, on average, the more active the overall Atlantic basin hurricane season is, the greater the probability of U.S. hurricane landfall.

Table 8: NTC activity in any year consists of the seasonal total of the following six parameters expressed in terms of their long-term averages. A season with 10 NS, 50 NSD, 6 H, 25 HD, 3 MH, and 5 MHD would then be the sum of the following ratios: $10/9.6 = 104$, $50/49.1 = 102$, $6/5.9 = 102$, $25/24.5 = 102$, $3/2.3 = 130$, $5/5.0 = 100$, divided by six, yielding an NTC of 107.

1950-2000 Average	
1) Named Storms (NS)	9.6
2) Named Storm Days (NSD)	49.1
3) Hurricanes (H)	5.9
4) Hurricane Days (HD)	24.5
5) Major Hurricanes (MH)	2.3
6) Major Hurricane Days (MHD)	5.0

Table 9 lists strike probabilities for the 2009 hurricane season for different TC categories for the entire U.S. coastline, the Gulf Coast and the East Coast including the Florida peninsula. We are also debuting probabilities for various islands and landmasses in the Caribbean and in Central America with this early June prediction. We have included the probability of a tropical cyclone of various categories tracking into the Caribbean (defined as 10-20°N, 60-88°W) in Table 9. The mean annual probability of one or more landfalling systems is given in parentheses. Note that Atlantic basin NTC activity in 2009 is expected to be slightly below its long-term average of 100, and therefore, landfall probabilities are slightly below their long-term average.

Please visit the Landfalling Probability Webpage at <http://www.e-transit.org/hurricane> for landfall probabilities for 11 U.S. coastal regions and 205 coastal and near-coastal counties from Brownsville, Texas to Eastport, Maine. We now have included probabilities of named storms, hurricanes and major hurricanes tracking within 50 and 100 miles of various islands and landmasses in the Caribbean and Central America.

Table 9: Estimated probability (expressed in percent) of one or more landfalling tropical storms (TS), category 1-2 hurricanes (HUR), category 3-4-5 hurricanes, total hurricanes and named storms along the entire U.S. coastline, along the Gulf Coast (Regions 1-4), and along the Florida Peninsula and the East Coast (Regions 5-11) for 2009. Probabilities of a tropical storm, hurricane and major hurricane tracking into the Caribbean are also provided. The long-term mean annual probability of one or more landfalling systems during the last 100 years is given in parentheses.

Region	TS	Category 1-2 HUR	Category 3-4-5 HUR	All HUR	Named Storms
Entire U.S. (Regions 1-11)	76% (79%)	64% (68%)	48% (52%)	81% (84%)	95% (97%)
Gulf Coast (Regions 1-4)	55% (59%)	39% (42%)	28% (30%)	56% (60%)	80% (83%)
Florida plus East Coast (Regions 5-11)	47% (50%)	41% (44%)	28% (31%)	57% (61%)	77% (81%)
Caribbean (10-20°N, 60-88°W)	79% (82%)	53% (57%)	39% (42%)	71% (75%)	94% (96%)

9 Has Global Warming Been Responsible for the Recent Large Upswing (Since 1995) in Atlantic Basin Major Hurricanes and U.S. Landfall?

The U.S. landfall of major hurricanes Dennis, Katrina, Rita and Wilma in 2005 and the four Southeast landfalling hurricanes of 2004 (Charley, Frances, Ivan and Jeanne) raised questions about the possible role that global warming played in these two unusually destructive seasons. In addition, three Category 2 hurricanes (Dolly, Gustav and Ike) pummeled the Gulf Coast last year causing considerable devastation.

The global warming arguments have been given much attention by many media references to recent papers claiming to show such a linkage. Despite the global warming

of the sea surface that has taken place over the last three decades, the global numbers of hurricanes and their intensity have not shown increases in recent years except for the Atlantic (Klotzbach 2006).

The Atlantic has seen a very large increase in major hurricanes during the 14-year period of 1995-2008 (average 3.9 per year) in comparison to the prior 25-year period of 1970-1994 (average 1.5 per year). This large increase in Atlantic major hurricanes is primarily a result of the multi-decadal increase in the Atlantic Ocean thermohaline circulation (THC) that is not directly related to global sea surface temperatures or CO₂ increases. Changes in ocean salinity are believed to be the driving mechanism. These multi-decadal changes have also been termed the Atlantic Multidecadal Oscillation (AMO). The AMO is the Atlantic component of the global ocean Meridional Overturning Circulation (MOC).

Although global surface temperatures have increased over the last century and over the last 30 years, there is no reliable data available to indicate increased hurricane frequency or intensity in any of the globe's other tropical cyclone basins.

In a global warming or global cooling world, the atmosphere's upper air temperatures will warm or cool in unison with the sea surface temperatures. Vertical lapse rates will not be significantly altered. We have no plausible physical reasons for believing that Atlantic hurricane frequency or intensity will change significantly if global ocean temperatures were to continue to rise. For instance, in the quarter-century period from 1945-1969 when the globe was undergoing a weak cooling trend, the Atlantic basin experienced 80 major (Cat 3-4-5) hurricanes and 201 major hurricane days. By contrast, in a similar 25-year period from 1970-1994 when the globe was undergoing a general warming trend, there were only 38 major hurricanes (48% as many) and 63 major hurricane days (31% as many) (Figure 10). Atlantic sea surface temperatures and hurricane activity do not necessarily follow global mean temperature trends.




Figure 10: Tracks of major (Category 3-4-5) hurricanes during the 25-year period of 1945-1969 when the globe was undergoing a weak cooling versus the 25-year period of 1970-1994 when the globe was undergoing a modest warming. CO₂ amounts in the later period were approximately 18 percent higher than in the earlier period. Major Atlantic hurricane activity was only about one-third as frequent during the latter period despite warmer global temperatures and higher CO₂ amounts.

The most reliable long-period hurricane records we have are the measurements of US landfalling tropical cyclones since 1899 (Table 10). Although global mean ocean and Atlantic sea surface temperatures have increased by about 0.4°C between these two 55-year periods (1899-1953 compared with 1954-2008), the frequency of US landfall numbers actually shows a slight downward trend for the later period. This downward trend is particularly noticeable for the US East Coast and Florida Peninsula where the difference in landfall of major (Category 3-4-5) hurricanes between the 43-year period of 1923-1965 (24 landfall events) and the 43-year period of 1966-2008 (7 landfall events) was especially large (Figure 11). For the entire United States coastline, 38 major hurricanes made landfall during the earlier 43-year period (1923-1965) compared with only 26 for the latter 43-year period (1966-2008). This occurred despite the fact that CO₂ averaged approximately 365 ppm during the latter period compared with 310 ppm during the earlier period.

Table 10: U.S. landfalling tropical cyclones by intensity during two 55-year periods.

YEARS	Named Storms	Hurricanes	Intense Hurricanes (Cat 3-4-5)	Global Temperature Increase
1899-1953 (55 years)	207	111	42	+0.4°C
1954-2008 (55 years)	188	95	39	

We should not read too much into the two hurricane seasons of 2004-2005. The activity of these two years was unusual but well within the natural bounds of hurricane variation.

What made the 2004-2005 and 2008 seasons so destructive was not the high frequency of major hurricanes but the high percentage of hurricanes that were steered over the US coastline. The US hurricane landfall events of these years were primarily a result of the favorable upper-air steering currents present during these years.




Figure 11: Contrast of tracks of East Coast and Florida Peninsula major landfalling hurricanes during the 43-year period of 1923-1965 versus the most recent 43-year period of 1966-2008.

Although 2005 had a record number of tropical cyclones (28 named storms), this should not be taken as an indication of something beyond natural processes. There have been several other years with comparable hurricane activity to 2005. For instance, 1933

had 21 named storms in a year when there was no satellite or aircraft data. Records of 1933 show all 21 named storm had tracks west of 60°W where surface observations were more plentiful. If we eliminate all the named storms of 2005 whose tracks were entirely east of 60°W and therefore may have been missed given the technology available in 1933, we reduce the 2005 named storm total by seven (to 21) – the same number as was observed to occur in 1933.

Utilizing the National Hurricanes Center’s best track database of hurricane records back to 1875, six previous seasons had more hurricane days than the 2005 season. These years were 1878, 1893, 1926, 1933, 1950 and 1995. Also, five prior seasons (1893, 1926, 1950, 1961 and 2004) had more major hurricane days. Although the 2005 hurricane season was certainly one of the most active on record, it was not as much of an outlier as many have indicated.

The active hurricane season in 2008 lends further support to the belief that the Atlantic basin remains in an active hurricane cycle associated with a strong thermohaline circulation and an active phase of the Atlantic Multidecadal Oscillation (AMO). This active cycle is expected to continue for another decade or two at which time we should enter a quieter Atlantic major hurricane period like we experienced during the quarter-century periods of 1970-1994 and 1901-1925. Atlantic hurricanes go through multi-decadal cycles. Cycles in Atlantic major hurricanes have been observationally traced back to the mid-19th century, and changes in the AMO have been inferred from Greenland paleo ice-core temperature measurements going back thousand of years.

10 Anticipated Large Increase in US Hurricane Destruction

The large increase in the hurricane-spawned destruction that occurred in 2004, 2005 and 2008 has not surprised us. We have been anticipating a great upsurge in hurricane destruction for many years as illustrated by the statements we have made in previous seasonal forecast reports such as:

“...major increases in hurricane-spawned coastal destruction are inevitable.”
(April 1989)

“A new era of major hurricane activity appears to have begun.... As a consequence of the exploding U.S. and Caribbean coastal populations during the last 25-30 years, we will begin to see a large upturn in hurricane-spawned destruction – likely higher than anything previous experienced.” (June 1997)

“We must expect a great increase in landfalling major hurricanes in the coming decades. With exploding southeast coastal populations, we must also prepare for levels of hurricane damage never before experienced.” (April 2001)

“If the future is like the past, it is highly likely that very active hurricane seasons will again emerge during the next few years, and the prospects for very large U.S. and Caribbean increases in hurricane damage over the next few decades remains high. We

should indeed see future hurricane damage much greater than anything in the past.” (May 2002)

“Regardless of whether a major hurricane makes landfall this year, it is inevitable that we will see hurricane-spawned destruction in coming years on a scale many, many times greater than what we have seen in the past.” (May 2003)

These projections of increased U.S. hurricane destruction were made with our anticipation that the Atlantic thermohaline circulation (THC) (which had been very weak from the late-1960s to the mid-1990s) would be changing to a stronger mode making for a large increase in Atlantic basin major hurricane activity. The THC has become much stronger since 1995. **These projections were made with no consideration given to rising levels of atmospheric CO₂.**

We were very fortunate during the early part of this strong THC period in that only 3 of 32 major hurricanes that formed in the Atlantic between 1995-2003 made U.S. landfall. The long-term average is that approximately 1 in 3.5 major hurricanes that forms in the Atlantic makes U.S. landfall. This luck failed to hold beginning with the 2004 hurricane season.

11 Forthcoming Updated Forecasts of 2009 Hurricane Activity

We will be issuing a final seasonal updates of our 2009 Atlantic basin hurricane forecast on **Tuesday 4 August**. We are planning on issuing experimental 15-day forecasts approximately every two weeks during the August-October period. These 15-day forecasts will supersede the August, September and October monthly forecasts. A verification and discussion of all 2009 forecasts will be issued in late November 2009. Our first seasonal hurricane forecast for the 2010 hurricane season will be issued in early December 2009. All of these forecasts will be available on the web at: <http://hurricane.atmos.colostate.edu/Forecasts>.

12 Acknowledgments

Besides the individuals named on page 5, there have been a number of other meteorologists that have furnished us with data and given valuable assessments of the current state of global atmospheric and oceanic conditions. These include Brian McNoldy, Arthur Douglas, Ray Zehr, Mark DeMaria, Todd Kimberlain, Paul Roundy and Amato Evan. In addition, Barbara Brumit and Amie Hedstrom have provided excellent manuscript, graphical and data analysis and assistance over a number of years. We have profited over the years from many in-depth discussions with most of the current and past NHC hurricane forecasters. The second author would further like to acknowledge the encouragement he has received for this type of forecasting research application from Neil Frank, Robert Sheets, Robert Burpee, Jerry Jarrell, and Max Mayfield, former directors of the National Hurricane Center (NHC). Uma Shama, Larry Harman and Daniel Fitch of Bridgewater State College, MA have provided assistance and technical support in the development of our Landfalling Hurricane Probability

Webpage. We also thank Bill Bailey of the Insurance Information Institute for his sage advice and encouragement.

The financial backing for the issuing and verification of these forecasts has been supported in part by the National Science Foundation. We also thank the GeoGraphics Laboratory at Bridgewater State College for their assistance in developing the Landfalling Hurricane Probability Webpage.

13 Citations and Additional Reading

- Blake, E. S., 2002: Prediction of August Atlantic basin hurricane activity. Dept. of Atmos. Sci. Paper No. 719, Colo. State Univ., Ft. Collins, CO, 80 pp.
- Blake, E. S. and W. M. Gray, 2004: Prediction of August Atlantic basin hurricane activity. *Wea. Forecasting*, 19, 1044-1060.
- Chiang, J. C. H. and D. J. Vimont, 2004: Analogous Pacific and Atlantic meridional modes of tropical atmosphere-ocean variability. *J. Climate*, 17, 4143-4158.
- DeMaria, M., J. A. Knaff and B. H. Connell, 2001: A tropical cyclone genesis parameter for the tropical Atlantic. *Wea. Forecasting*, 16, 219-233.
- Elsner, J. B., G. S. Lehmiller, and T. B. Kimberlain, 1996: Objective classification of Atlantic hurricanes. *J. Climate*, 9, 2880-2889.
- Evan, A. T., J. Dunion, J. A. Foley, A. K. Heidinger, and C. S. Velden, 2006: New evidence for a relationship between Atlantic tropical cyclone activity and African dust outbreaks, *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL026408.
- Goldenberg, S. B., C. W. Landsea, A. M. Mestas-Nunez, and W. M. Gray, 2001: The recent increase in Atlantic hurricane activity: Causes and Implications. *Science*, 293, 474-479.
- Goldenberg, S. B. and L. J. Shapiro, 1996: Physical mechanisms for the association of El Niño and West African rainfall with Atlantic major hurricane activity. *J. Climate*, 1169-1187.
- Gray, W. M., 1984a: Atlantic seasonal hurricane frequency: Part I: El Niño and 30 mb quasi-biennial oscillation influences. *Mon. Wea. Rev.*, 112, 1649-1668.
- Gray, W. M., 1984b: Atlantic seasonal hurricane frequency: Part II: Forecasting its variability. *Mon. Wea. Rev.*, 112, 1669-1683.
- Gray, W. M., 1990: Strong association between West African rainfall and US landfall of intense hurricanes. *Science*, 249, 1251-1256.
- Gray, W. M., and P. J. Klotzbach, 2003 and 2004: Forecasts of Atlantic seasonal and monthly hurricane activity and US landfall strike probability. Available online at <http://hurricane.atmos.colostate.edu>
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1992: Predicting Atlantic seasonal hurricane activity 6-11 months in advance. *Wea. Forecasting*, 7, 440-455.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1993: Predicting Atlantic basin seasonal tropical cyclone activity by 1 August. *Wea. Forecasting*, 8, 73-86.

- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1994a: Predicting Atlantic basin seasonal tropical cyclone activity by 1 June. *Wea. Forecasting*, 9, 103-115.
- Gray, W. M., J. D. Sheaffer and C. W. Landsea, 1996: Climate trends associated with multi-decadal variability of intense Atlantic hurricane activity. Chapter 2 in "Hurricanes, Climatic Change and Socioeconomic Impacts: A Current Perspective", H. F. Diaz and R. S. Pulwarty, Eds., Westview Press, 49 pp.
- Gray, W. M., 1998: Atlantic ocean influences on multi-decadal variations in El Niño frequency and intensity. Ninth Conference on Interaction of the Sea and Atmosphere, 78th AMS Annual Meeting, 11-16 January, Phoenix, AZ, 5 pp.
- Henderson-Sellers, A., H. Zhang, G. Berz, K. Emanuel, W. Gray, C. Landsea, G. Holland, J. Lighthill, S-L. Shieh, P. Webster, and K. McGuffie, 1998: Tropical cyclones and global climate change: A post-IPCC assessment. *Bull. Amer. Meteor. Soc.*, 79, 19-38.
- Klotzbach, P. J., 2002: Forecasting September Atlantic basin tropical cyclone activity at zero and one-month lead times. Dept. of Atmos. Sci. Paper No. 723, Colo. State Univ., Ft. Collins, CO, 91 pp.
- Klotzbach, P. J., 2006: Trends in global tropical cyclone activity over the past twenty years (1986-2005). *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL025881.
- Klotzbach, P. J., 2007: Revised prediction of seasonal Atlantic basin tropical cyclone activity from 1 August. *Wea. and Forecasting*, 22, 937-949.
- Klotzbach, P. J. and W. M. Gray, 2003: Forecasting September Atlantic basin tropical cyclone activity. *Wea. and Forecasting*, 18, 1109-1128.
- Klotzbach, P. J. and W. M. Gray, 2004: Updated 6-11 month prediction of Atlantic basin seasonal hurricane activity. *Wea. and Forecasting*, 19, 917-934.
- Klotzbach, P. J. and W. M. Gray, 2006: Causes of the unusually destructive 2004 Atlantic basin hurricane season. *Bull. Amer. Meteor. Soc.*, 87, 1325-1333.
- Klotzbach, P. J. and W. M. Gray, 2009: Twenty-five years of Atlantic basin seasonal hurricane forecasts. *Geophys. Res. Lett.*, 36, L09711, doi:10.1029/2009GL037580.
- Knaff, J. A., 1997: Implications of summertime sea level pressure anomalies. *J. Climate*, 10, 789-804.
- Knaff, J. A., 1998: Predicting summertime Caribbean sea level pressure. *Wea. and Forecasting*, 13, 740-752.
- Kossin, J. P., and D. J. Vimont, 2007: A more general framework for understanding Atlantic hurricane variability and trends. *Bull. Amer. Meteor. Soc.*, 88, 1767-1781.
- Landsea, C. W., 1991: West African monsoonal rainfall and intense hurricane associations. Dept. of Atmos. Sci. Paper, Colo. State Univ., Ft. Collins, CO, 272 pp.
- Landsea, C. W., 1993: A climatology of intense (or major) Atlantic hurricanes. *Mon. Wea. Rev.*, 121, 1703-1713.
- Landsea, C. W., 2007: Counting Atlantic tropical cyclones back to 1900. *EOS*, 88, 197, 202.
- Landsea, C. W. and W. M. Gray, 1992: The strong association between Western Sahel monsoon rainfall and intense Atlantic hurricanes. *J. Climate*, 5, 435-453.

- Landsea, C. W., W. M. Gray, P. W. Mielke, Jr., and K. J. Berry, 1992: Long-term variations of Western Sahelian monsoon rainfall and intense U.S. landfalling hurricanes. *J. Climate*, 5, 1528-1534.
- Landsea, C. W., W. M. Gray, K. J. Berry and P. W. Mielke, Jr., 1996: June to September rainfall in the African Sahel: A seasonal forecast for 1996. 4 pp.
- Landsea, C. W., N. Nicholls, W.M. Gray, and L.A. Avila, 1996: Downward trends in the frequency of intense Atlantic hurricanes during the past five decades. *Geo. Res. Letters*, 23, 1697-1700.
- Landsea, C. W., R. A. Pielke, Jr., A. M. Mestas-Nunez, and J. A. Knaff, 1999: Atlantic basin hurricanes: Indices of climatic changes. *Climatic Changes*, 42, 89-129.
- Landsea, C.W. et al., 2005: Atlantic hurricane database re-analysis project. Available online at http://www.aoml.noaa.gov/hrd/data_sub/re_anal.html
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1996: Artificial skill and validation in meteorological forecasting. *Wea. Forecasting*, 11, 153-169.
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1997: A single sample estimate of shrinkage in meteorological forecasting. *Wea. Forecasting*, 12, 847-858.
- Pielke, Jr. R. A., and C. W. Landsea, 1998: Normalized Atlantic hurricane damage, 1925-1995. *Wea. Forecasting*, 13, 621-631.
- Rasmusson, E. M. and T. H. Carpenter, 1982: Variations in tropical sea-surface temperature and surface wind fields associated with the Southern Oscillation/El Niño. *Mon. Wea. Rev.*, 110, 354-384.
- Seseske, S. A., 2004: Forecasting summer/fall El Niño-Southern Oscillation events at 6-11 month lead times. Dept. of Atmos. Sci. Paper No. 749, Colo. State Univ., Ft. Collins, CO, 104 pp.
- Vimont, D. J., and J. P. Kossin, 2007: The Atlantic meridional mode and hurricane activity. *Geophys. Res. Lett.*, 34, L07709, doi:10.1029/2007GL029683.

14 Verification of Previous Forecasts

Table 11: Summary verification of the authors' five previous years of seasonal forecasts for Atlantic TC activity between 2004-2008.

2004	5 Dec. 2003	Update 2 April	Update 28 May	Update 6 August	Obs.
Hurricanes	7	8	8	7	9
Named Storms	13	14	14	13	14
Hurricane Days	30	35	35	30	46
Named Storm Days	55	60	60	55	90
Intense Hurricanes	3	3	3	3	6
Intense Hurricane Days	6	8	8	6	22
Net Tropical Cyclone Activity	125	145	145	125	229

2005	3 Dec. 2004	Update 1 April	Update 31 May	Update 5 August	Obs.
Hurricanes	6	7	8	10	14
Named Storms	11	13	15	20	26
Hurricane Days	25	35	45	55	48
Named Storm Days	55	65	75	95	116
Intense Hurricanes	3	3	4	6	7
Intense Hurricane Days	6	7	11	18	16.75
Net Tropical Cyclone Activity	115	135	170	235	263

2006	6 Dec. 2005	Update 4 April	Update 31 May	Update 3 August	Obs.
Hurricanes	9	9	9	7	5
Named Storms	17	17	17	15	10
Hurricane Days	45	45	45	35	20
Named Storm Days	85	85	85	75	50
Intense Hurricanes	5	5	5	3	2
Intense Hurricane Days	13	13	13	8	3
Net Tropical Cyclone Activity	195	195	195	140	85

2007	8 Dec. 2006	Update 3 April	Update 31 May	Update 3 August	Obs.
Hurricanes	7	9	9	8	6
Named Storms	14	17	17	15	15
Hurricane Days	35	40	40	35	11.25
Named Storm Days	70	85	85	75	34.50
Intense Hurricanes	3	5	5	4	2
Intense Hurricane Days	8	11	11	10	5.75
Net Tropical Cyclone Activity	140	185	185	160	97

2008	7 Dec. 2007	Update 9 April	Update 3 June	Update 5 August	Obs.
Hurricanes	7	8	8	9	8
Named Storms	13	15	15	17	16
Hurricane Days	30	40	40	45	30.50
Named Storm Days	60	80	80	90	88.25
Intense Hurricanes	3	4	4	5	5
Intense Hurricane Days	6	9	9	11	7.50
Net Tropical Cyclone Activity	125	160	160	190	162