

EXTENDED RANGE FORECAST OF ATLANTIC SEASONAL HURRICANE ACTIVITY AND LANDFALL STRIKE PROBABILITY FOR 2021

We have maintained our above-average forecast for the 2021 Atlantic basin hurricane season. Current neutral ENSO conditions are anticipated to persist for the next several months. While sea surface temperatures averaged across portions of the tropical Atlantic are near to slightly below normal, subtropical North Atlantic sea surface temperatures are much warmer than average. We anticipate an above-normal probability for major hurricanes making landfall along the continental United States coastline and in the Caribbean. As is the case with all hurricane seasons, coastal residents are reminded that it only takes one hurricane making landfall to make it an active season for them. They should prepare the same for every season, regardless of how much activity is predicted.

(as of 3 June 2021)

By Philip J. Klotzbach¹, Michael M. Bell², and Jhordanne Jones³

In Memory of William M. Gray⁴

This discussion as well as past forecasts and verifications are available online at <http://tropical.colostate.edu>

Jennifer Dimas, Colorado State University media representative, is coordinating media inquiries into this verification. She can be reached at 970-491-1543 or Jennifer.Dimas@colostate.edu

Department of Atmospheric Science
Colorado State University
Fort Collins, CO 80523

Project Sponsors:



¹ Research Scientist

² Associate Professor

³ Graduate Research Assistant

⁴ Professor Emeritus

ATLANTIC BASIN SEASONAL HURRICANE FORECAST FOR 2021

| Forecast Parameter and 1991-2020 Average (in parentheses) | Issue Date 8 April 2021 | Issue Date 3 June 2021 | Observed Activity Through June 2 2021 | Total Seasonal Forecast (Includes Ana*) |
|--|-------------------------------|------------------------------|---|--|
| Named Storms (14.4) | 17 | 17 | 1 | 18 |
| Named Storm Days (69.4) | 80 | 78.25 | 1.75 | 80 |
| Hurricanes (7.2) | 8 | 8 | 0 | 8 |
| Hurricane Days (27.0) | 35 | 35 | 0 | 35 |
| Major Hurricanes (3.2) | 4 | 4 | 0 | 4 |
| Major Hurricane Days (7.4) | 9 | 9 | 0 | 9 |
| Accumulated Cyclone Energy Index (123) | 150 | 149 | 1 | 150 |
| Net Tropical Cyclone Activity (135%) | 160 | 158 | 2 | 160 |

*Ana formed prior to the start of the Atlantic hurricane season on June 1st.

PROBABILITIES FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE LANDFALL ON EACH OF THE FOLLOWING COASTAL AREAS:

- 1) Entire continental U.S. coastline - 69% (average for last century is 52%)
- 2) U.S. East Coast Including Peninsula Florida - 45% (average for last century is 31%)
- 3) Gulf Coast from the Florida Panhandle westward to Brownsville - 44% (average for last century is 30%)

PROBABILITY FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE TRACKING INTO THE CARIBBEAN (10-20°N, 88-60°W)

- 1) 58% (average for last century is 42%)

ABSTRACT

Information obtained through May 2021 indicates that the 2021 Atlantic hurricane season will have activity slightly above the 1991-2020 average. We estimate that 2021 will have 8 hurricanes (average is 7.2), 18 named storms (average is 14.4), 80 named storm days (average is 69.4), 35 hurricane days (average is 27.0), 4 major (Category 3-4-5) hurricanes (average is 3.2) and 9 major hurricane days (average is 7.4). The probability of U.S. major hurricane landfall is estimated to be about 135 percent of the long-period average. We expect Atlantic basin Accumulated Cyclone Energy (ACE) and Net Tropical Cyclone (NTC) activity in 2021 to be approximately 120 percent of their long-term averages.

This forecast is based on an extended-range early June statistical prediction scheme that was developed using 39 years of past data. Analog predictors are also utilized. We are also including statistical/dynamical models based off data from both the ECMWF SEAS5 model and the Met Office GloSea6 model as two additional forecast guidance tools. The statistical model, the two statistical/dynamical models and the analog model all call for an above-average Atlantic hurricane season. We also present probabilities of exceedance for hurricanes and Accumulated Cyclone Energy to give interested readers a better idea of the uncertainty associated with these forecasts.

The tropical Pacific is currently characterized by neutral ENSO conditions, with anomalous eastern and central tropical Pacific warming over the past several months putting an end to this past winter's La Niña event. We anticipate that neutral ENSO conditions are the most likely scenario for the peak of this year's Atlantic hurricane season, and it seems unlikely that El Niño conditions will develop over the next few months. El Niño typically reduces Atlantic hurricane activity through increases in vertical wind shear.

The tropical Atlantic currently has near-normal sea surface temperatures, while most of the subtropical North Atlantic is warmer than normal. A warmer subtropical North Atlantic in the late spring typically correlates with a weaker subtropical high that then leads to anomalous warming of the tropical Atlantic by the peak of the Atlantic hurricane season.

Coastal residents are reminded that it only takes one hurricane making landfall to make it an active season for them, and they need to prepare the same for every season, regardless of how much activity is predicted.

The early June forecast has moderate long-term skill when evaluated in hindcast mode. The skill of CSU's forecast updates increases as the peak of the Atlantic hurricane season approaches.

Why issue extended-range forecasts for seasonal hurricane activity?

We are frequently asked this question. Our answer is that it is possible to say something about the probability of the coming year's hurricane activity which is superior to climatology. The Atlantic basin has the largest year-to-year variability of any of the global tropical cyclone basins. People are curious to know how active the upcoming season is likely to be, particularly if you can show hindcast skill improvement over climatology for many past years.

Everyone should realize that it is impossible to precisely predict this season's hurricane activity in early June. There is, however, much curiosity as to how global ocean and atmosphere features are presently arranged as regards to the probability of an active or inactive hurricane season for the coming year. Our early June statistical and statistical/dynamical hybrid models show strong evidence on ~25-40 years of data that significant improvement over a climatological forecast can be attained. We would never issue a seasonal hurricane forecast unless we had models developed over a long hindcast period which showed skill. We also now include probabilities of exceedance to provide improved quantification of the uncertainty associated with these predictions.

We issue these forecasts to satisfy the curiosity of the general public and to bring attention to the hurricane problem. There is a general interest in knowing what the odds are for an active or inactive season. One must remember that our forecasts are based on the premise that those global oceanic and atmospheric conditions which preceded comparatively active or inactive hurricane seasons in the past provide meaningful information about similar trends in future seasons.

It is also important that the reader appreciate that these seasonal forecasts are based on statistical and dynamical models which will fail in some years. Moreover, these forecasts do not specifically predict where within the Atlantic basin these storms will strike. The probability of landfall for any one location along the coast is very low and reflects the fact that, in any one season, most U.S. coastal areas will not feel the effects of a hurricane no matter how active the individual season is.

Acknowledgment

These seasonal forecasts were developed by the late Dr. William Gray, who was lead author on these predictions for over 20 years and continued as a co-author until his death in 2016. In addition to pioneering seasonal Atlantic hurricane prediction, he conducted groundbreaking research in a wide variety of other topics including hurricane genesis, hurricane structure and cumulus convection. His investments in both time and energy to these forecasts cannot be acknowledged enough.

We are grateful for support from Ironshore Insurance, the Insurance Information Institute, Weatherboy and Evex. We acknowledge a grant from the G. Unger Vetlesen Foundation for additional financial support.

Colorado State University's seasonal hurricane forecasts have benefited greatly from a number of individuals that were former graduate students of William Gray. Among these former project members are Chris Landsea, John Knaff and Eric Blake. We would like to acknowledge assistance from Louis-Philippe Caron and the data team at the Barcelona Supercomputing Centre for providing data and insight on the statistical/dynamical models. We have also benefited from meteorological discussions with Carl Schreck, Louis-Philippe Caron, Brian McNoldy, Paul Roundy, Jason Dunion, Peng Xian and Amato Evan over the past few years.

DEFINITIONS AND ACRONYMS

Accumulated Cyclone Energy (ACE) - A measure of a named storm's potential for wind and storm surge destruction defined as the sum of the square of a named storm's maximum wind speed (in 10^4 knots²) for each 6-hour period of its existence. The 1981–2010 average value of this parameter is 106 for the Atlantic basin.

Atlantic Multi-Decadal Oscillation (AMO) – A mode of natural variability that occurs in the North Atlantic Ocean and evidencing itself in fluctuations in sea surface temperature and sea level pressure fields. The AMO is likely related to fluctuations in the strength of the oceanic thermohaline circulation. Although several definitions of the AMO are currently used in the literature, we define the AMO based on North Atlantic sea surface temperatures from 50-60°N, 50-10°W and sea level pressure from 0-50°N, 70-10°W.

Atlantic Basin – The area including the entire North Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico.

El Niño – A 12-18 month period during which anomalously warm sea surface temperatures occur in the eastern half of the equatorial Pacific. Moderate or strong El Niño events occur irregularly, about once every 3-7 years on average.

Hurricane (H) - A tropical cyclone with sustained low-level winds of 74 miles per hour (33 ms^{-1} or 64 knots) or greater.

Hurricane Day (HD) - A measure of hurricane activity, one unit of which occurs as four 6-hour periods during which a tropical cyclone is observed or is estimated to have hurricane-force winds.

Indian Ocean Dipole (IOD) - An irregular oscillation of sea surface temperatures between the western and eastern tropical Indian Ocean. A positive phase of the IOD occurs when the western Indian Ocean is anomalously warm compared with the eastern Indian Ocean.

Madden Julian Oscillation (MJO) – A globally propagating mode of tropical atmospheric intra-seasonal variability. The wave tends to propagate eastward at approximately 5 ms^{-1} , circling the globe in roughly 30-60 days.

Main Development Region (MDR) – An area in the tropical Atlantic where a majority of major hurricanes form, which we define as 7.5-22.5°N, 75-20°W.

Major Hurricane (MH) - A hurricane which reaches a sustained low-level wind of at least 111 mph (96 knots or 50 ms^{-1}) at some point in its lifetime. This constitutes a category 3 or higher on the Saffir/Simpson scale.

Major Hurricane Day (MHD) - Four 6-hour periods during which a hurricane has an intensity of Saffir/Simpson category 3 or higher.

Multivariate ENSO Index (MEI) – An index defining ENSO that takes into account tropical Pacific sea surface temperatures, sea level pressures, zonal and meridional winds and cloudiness.

Named Storm (NS) - A hurricane, a tropical storm or a sub-tropical storm.

Named Storm Day (NSD) - As in HD but for four 6-hour periods during which a tropical or sub-tropical cyclone is observed (or is estimated) to have attained tropical storm-force winds.

Net Tropical Cyclone (NTC) Activity – Average seasonal percentage mean of NS, NSD, H, HD, MH, MHD. Gives overall indication of Atlantic basin seasonal hurricane activity. The 1950-2000 average value of this parameter is 100.

Proxy – An approximation or a substitution for a physical process that cannot be directly measured.

Saffir/Simpson Hurricane Wind Scale – A measurement scale ranging from 1 to 5 of hurricane wind intensity. One is a weak hurricane; whereas, five is the most intense hurricane.

Southern Oscillation Index (SOI) – A normalized measure of the surface pressure difference between Tahiti and Darwin. Low values typically indicate El Niño conditions.

Standard Deviation (SD) – A measure used to quantify the variation in a dataset.

Sea Surface Temperature Anomaly – SSTA

Thermohaline Circulation (THC) – A large-scale circulation in the Atlantic Ocean that is driven by fluctuations in salinity and temperature. When the THC is stronger than normal, the AMO tends to be in its warm (or positive) phase, and more Atlantic hurricanes typically form.

Tropical Cyclone (TC) - A large-scale circular flow occurring within the tropics and subtropics which has its strongest winds at low levels; including hurricanes, tropical storms and other weaker rotating vortices.

Tropical North Atlantic (TNA) index – A measure of sea surface temperatures in the area from 5.5-23.5°N, 57.5-15°W.

Tropical Storm (TS) - A tropical cyclone with maximum sustained winds between 39 mph (18 ms^{-1} or 34 knots) and 73 mph (32 ms^{-1} or 63 knots).

Vertical Wind Shear – The difference in horizontal wind between 200 hPa (approximately 40000 feet or 12 km) and 850 hPa (approximately 5000 feet or 1.6 km).

1 knot = 1.15 miles per hour = 0.515 meters per second

1 Introduction

This is the 38th year in which the CSU Tropical Meteorology Project has made forecasts of the upcoming season's Atlantic basin hurricane activity. Our research team has shown that a sizable portion of the year-to-year variability of Atlantic tropical cyclone (TC) activity can be hindcast with skill exceeding climatology. This year's June forecast is based on a statistical methodology as well as output from two statistical/dynamical models calculated from the SEAS5 climate model from the European Centre for Medium Range Weather Forecasts (ECMWF) and the GloSea6 model from the UK Met Office. These models show skill on 25-40 years of historical data, depending on the particular forecast technique. We also select analog seasons, based primarily on conditions we anticipate for the peak of the Atlantic hurricane season. Qualitative adjustments are added to accommodate additional processes which may not be explicitly represented by these analyses. These evolving forecast techniques are based on a variety of climate-related global and regional predictors previously shown to be related to the forthcoming seasonal Atlantic basin TC activity and landfall probability. We believe that seasonal forecasts must be based on methods that show significant hindcast skill in application to long periods of prior data. It is only through hindcast skill that one can demonstrate that seasonal forecast skill is possible. This is a valid methodology provided that the atmosphere continues to behave in the future as it has in the past.

The best predictors do not necessarily have the best individual correlations with hurricane activity. The best forecast parameters are those that explain the portion of the variance of seasonal hurricane activity that are not associated with the other forecast variables. It is possible for an important hurricane forecast parameter to show little direct relationship to a predictand by itself but to have an important influence when included with a set of 2-3 other predictors.

A direct correlation of a forecast parameter may not be the best measure of the importance of this predictor to the skill of a 3-4 parameter forecast model. This is the nature of the seasonal or climate forecast problem where one is dealing with a very complicated atmospheric-oceanic system that is highly non-linear. There is a maze of changing physical linkages between the many variables. These linkages can undergo unknown changes from weekly to decadal time scales. It is impossible to understand how all of these processes interact with each other. But, it is still possible to develop a reliable statistical forecast scheme which incorporates a number of the climate system's non-linear interactions. Any seasonal or climate forecast scheme should show significant hindcast skill before it is used in real-time forecasts.

2 June Forecast Methodology

2.1 June Statistical Forecast Scheme

We are continuing to use the same June statistical forecast scheme that we debuted last year. The model uses the ECMWF Reanalysis 5 (ERA5) (Hersbach et al. 2020) as well as NOAA Optimum Interpolation (OI) SST (Reynolds et al. 2002). The ERA5 reanalysis currently extends from 1979 to present with a preliminary backward extension now available back to 1950. A benefit of the ERA5 reanalysis is that it is the first reanalysis from ECMWF that provides updates in near real-time, allowing for the same reanalysis product to be used for both hindcast model development as well as real-time analysis. The NOAA OISST (Reynolds et al. 2002) is available from 1982-present. This new model showed significant skill in cross-validated (e.g., leaving the year out of the developmental model that is being predicted) hindcasts of Accumulated Cyclone Energy (ACE) ($r = 0.72$) over the period from 1982-2019 and also successfully forecast the active 2020 Atlantic hurricane season (Figure 1).

Figure 2 displays the locations of each of our predictors, while Table 1 displays the individual linear correlations between each predictor and ACE over the 1982-2020 period. All predictors correlate significantly at the 5% level using a two-tailed Student's t-test and assuming that each year is independent of the prior year (e.g., the correlation between ACE in two consecutive years is very low). Table 2 displays the 2021 observed values for each of the four predictors in the statistical forecast scheme. Table 3 displays the statistical model output for the 2021 hurricane season. Three of the four predictors call for increased Atlantic hurricane activity in 2021.




Figure 1: Observed versus early June cross-validated hindcast values of ACE for the statistical model for 1982-2020.

June Forecast Predictors




Figure 2: Location of predictors for our early June extended-range statistical prediction for the 2021 hurricane season.

Table 1: Rank correlations between early June predictors and ACE over the period from 1982-2020.

| Predictor | Correlation w/ ACE |
|---|--------------------|
| 1) April-May SST (20°N-60°N, 40°W-15°W) (+) | 0.58 |
| 2) May SLP (20°N-40°N, 60°W-10°W) (-) | -0.31 |
| 3) May 10m U (5°S-5°N, 180°W-130°W) (-) | -0.51 |
| 4) April-May SST (35°S-15°S, 155°E-180°E) (+) | 0.61 |

Table 2: Listing of early June 2021 predictors for the 2021 hurricane season. A plus (+) means that positive deviations of the parameter are associated with increased hurricane activity, while a minus (-) means that negative deviations of the parameter are associated with increased hurricane activity. SD stands for standard deviation.

| Predictor | 2021 Forecast Value | Impact on 2021 TC Activity |
|---|---------------------|----------------------------|
| 1) April-May SST (20°N-60°N, 40°W-15°W) (+) | +1.6 SD | Enhance |
| 2) May SLP (20°N-40°N, 60°W-10°W) (-) | +1.7 SD | Suppress |
| 3) May 10m U (5°S-5°N, 180°W-130°W) (-) | -2.5 SD | Enhance |
| 4) April-May SST (35°S-15°S, 155°E-180°E) (+) | +0.6 SD | Enhance |

Table 3: Statistical model output for the 2021 Atlantic hurricane season and the final adjusted forecast.

| Forecast Parameter and 1991-2020 Average (in parentheses) | Statistical Forecast | Final Forecast |
|---|----------------------|----------------|
| Named Storms (14.4) | 16.6 | 18 |
| Named Storm Days (69.4) | 84.2 | 80 |
| Hurricanes (7.2) | 8.7 | 8 |
| Hurricane Days (27.0) | 35.1 | 35 |
| Major Hurricanes (3.2) | 4.1 | 4 |
| Major Hurricane Days (7.4) | 10.3 | 9 |
| Accumulated Cyclone Energy Index (123) | 158 | 150 |
| Net Tropical Cyclone Activity (135%) | 170 | 160 |

The locations and brief descriptions of the predictors for our early June statistical forecast are now discussed. It should be noted that all predictors correlate with physical features during August through October that are known to be favorable for elevated levels of hurricane activity. These factors are all generally related to August-October vertical wind shear in the Atlantic Main Development Region (MDR) from 10-20°N, 85-20°W as shown in Figure 3.




Figure 3: Vertical wind profile typically associated with (a) inactive Atlantic basin hurricane seasons and (b) active Atlantic basin hurricane seasons. Note that (b) has reduced levels of vertical wind shear.

For each of these predictors, we display a four-panel figure showing rank correlations between values of each predictor and August-October values of sea surface temperature (SST), sea level pressure (SLP), 200 hPa zonal wind, and 850 hPa zonal wind, respectively, during 1982-2019. In general, higher values of SST, lower values of SLP, anomalous westerlies at 850 hPa and anomalous easterlies at 200 hPa are associated with active Atlantic basin hurricane seasons. SST correlations are displayed using the NOAA OISST, while atmospheric field correlations are displayed using ERA5.

Predictor 1. April-May SST in the Subtropical and Mid-latitude eastern North Atlantic (+)

(20°N-60°N, 40°W-15°W)

Warmer-than-normal SSTs in the subtropical and mid-latitude eastern North Atlantic during April-May are associated with a weaker-than-normal subtropical high and reduced trade wind strength during the boreal spring (Knaff 1997). These warmer-than-normal SSTs in April-May are also correlated with weaker trade winds and weaker upper tropospheric westerly winds, lower-than-normal sea level pressure and above-normal SSTs in the tropical Atlantic during the following August-October period (Figure 4). All three of these August-October features are commonly associated with active Atlantic basin hurricane seasons, through reductions in vertical wind shear, increased vertical instability and increased mid-tropospheric moisture, respectively. Predictor 1 rank correlates quite strongly ($r = 0.58$) with ACE from 1982-2020. Predictor 1 also strongly correlates ($r = 0.65$) with August-October values of the Atlantic Meridional Mode (AMM) (Kossin and Vimont 2007) over the period from 1982-2020. The AMM has been shown to impact Atlantic hurricane activity through alterations in the position and intensity of the Atlantic Inter-Tropical Convergence Zone (ITCZ). Changes in the Atlantic ITCZ bring about changes in tropical Atlantic vertical and horizontal wind shear patterns and in tropical Atlantic SST patterns.

Predictor 2. May SLP in the Subtropical North Atlantic (-)

(20°N-40°N, 60°W-10°W)

Anomalously low pressure in the subtropical North Atlantic during May is associated with weaker trade winds and anomalous warming of the central and eastern tropical Atlantic during the boreal spring. While the anomalous warming signal in the tropical Atlantic SST decays by the peak of the Atlantic hurricane season (Figure 5), there remains a significant signal in sea level pressure and vertical wind shear, as evidenced by the anomalous westerlies at 850 hPa and the anomalous easterlies at 200 hPa in the Caribbean and western tropical Atlantic during August-October. Consequently, we observe that when the subtropical North Atlantic is characterized by lower pressure in May, the peak of the Atlantic hurricane season typically has reduced levels of vertical wind shear across the Main Development Region.

Predictor 3. May 10m U in the central and eastern tropical Pacific (-)

(5°S-5°N, 180°W-130°W)

Stronger-than-normal low-level winds during May in the central and eastern tropical Pacific are associated with enhanced upwelling which drives anomalous cooling in the central and eastern tropical Pacific, inhibiting the development of El Niño conditions. This relationship can be clearly demonstrated by a significant correlation between Predictor 3 with the August-October-averaged Oceanic Niño Index ($r = 0.51$). As would be expected given this significant correlation, Predictor 3 also correlates with reduced vertical wind shear during the peak of the Atlantic hurricane season, especially in the Caribbean and western tropical Atlantic, where ENSO typically has its strongest impacts (Figure 6).

Predictor 4. April-May SST in the Tropical and Subtropical Western South Pacific
(+)

(35°S-15°S, 155°E-180°E)

Anomalous warmth in the tropical and subtropical western South Pacific is associated with higher-than-normal pressure in the eastern tropical Pacific during the boreal spring. This anomalous pressure pattern results in a positive Southern Oscillation Index (SOI), both in April-May and in August-October. The correlation between the SOI in April-May and Predictor 4 is 0.31, while the correlation increases to 0.42 between Predictor 4 and the August-October-averaged SOI. In addition, the correlation between Predictor 4 and the August-October averaged ENSO Longitude Index is -0.40. All three of these correlations indicate that when April-May SSTs in the tropical and subtropical western South Pacific are warmer than normal, the Walker Circulation is typically stronger and is consequently associated with decreased vertical wind shear across the Caribbean and tropical Atlantic for the peak of the Atlantic hurricane season (Figure 7).

August-October Correlations w/ Predictor 1 (1982-2019)




Figure 4: Rank correlations between April-May SST in the subtropical and mid-latitude eastern North Atlantic (Predictor 1) and (panel a) August-October sea surface temperature, (panel b) August-October sea level pressure, (panel c) August-October 850 hPa zonal wind and (panel d) August-October 200 hPa zonal wind. All four of these parameter deviations in the tropical Atlantic are known to be favorable for enhanced hurricane activity.

August-October Correlations w/ Predictor 2 (1982-2019)




Figure 5: As in Figure 4 but for May SLP in the subtropical North Atlantic. The sign of Predictor 2 has been flipped for easy comparison with Figure 4.

August-October Correlations w/ Predictor 3 (1982-2019)




Figure 6: As in Figure 4 but for 10 meter zonal wind in the central and eastern equatorial Pacific Ocean. The sign of Predictor 3 has been flipped for easy comparison with Figure 4.

August-October Correlations w/ Predictor 4 (1982-2019)




Figure 7: As in Figure 4 but for April-May SST in the tropical and subtropical western South Pacific.

2.2 June Statistical/Dynamical Forecast Scheme

We developed a new statistical/dynamical hybrid forecast model scheme that we used for the first time in 2019. This model, developed in partnership with Louis-Philippe Caron and the data team at the Barcelona Supercomputing Centre, uses output from the ECMWF SEAS5 model to forecast the input to our early August statistical forecast model. The early August statistical forecast model shows the highest level of skill of any of our statistical models, since it is the model released just before the peak of the Atlantic hurricane season in September. ECMWF SEAS5 is able to forecast the large-scale fields that go into the early August statistical forecast model with considerable skill by March. We then use the forecasts of the individual parameters to forecast ACE for the 2021 season. All other predictands (e.g., named storms, major hurricanes) are calculated based on their historical relationships with ACE. It typically takes about two weeks after the initialization date to obtain SEAS5 output, so the results displayed here are from the model output from the 1 May forecast.

Figure 8 displays the parameters used in our early August statistical model, while Table 4 displays SEAS5's forecasts of these parameters for 2021 from a 1 May initialization date. Two of the three parameters call for above-normal activity, while the upper-level wind predictor over Africa calls for near-average activity. Given the strongly positive forecast temperatures in the subtropical northeastern Atlantic, the SEAS5 statistical/dynamical model is calling for an active season. Figure 9 displays cross-validated hindcasts for SEAS5 forecast of ACE from 1981-2018 as well as the real-time forecasts for 2019-2020, while Table 5 presents the forecast from SEAS5 for the 2021 Atlantic hurricane season.

Post-31 July Seasonal Forecast Predictors




Figure 8: Location of predictors for our statistical/dynamical extended-range statistical prediction for the 2021 hurricane season. This forecast uses either the ECMWF SEAS5 model or the UK Met Office GloSea6 model to predict July conditions in the three boxes displayed and uses those predictors to forecast ACE.

Table 4: Listing of predictions of July large-scale conditions from ECMWF SEAS5 output, initialized on 1 May. A plus (+) means that positive deviations of the parameter are associated with increased hurricane activity, while a minus (-) means that negative deviations of the parameter are associated with increased hurricane activity.

| Predictor | Values for 2021 Forecast | Effect on 2021 Hurricane Season |
|---|--------------------------|---------------------------------|
| 1) ECMWF Prediction of July Surface U (10-20°N, 90-40°W) (+) | +0.2 SD | Enhance |
| 2) ECMWF Prediction of July Skin Temperature (20-40°N, 35-15°W) (+) | +1.5 SD | Enhance |
| 3) ECMWF Prediction of July 200 hPa U (5-15°N, 0-40°E) (-) | 0.0 SD | Neutral |




Figure 9: Observed versus May cross-validated statistical/dynamical hindcast values of ACE for 1981-2018 as well as real-time forecasts from SEAS5 for 2019-2020.

Table 5: Statistical/dynamical model output from SEAS5 for the 2021 Atlantic hurricane season and the final adjusted forecast.

| Forecast Parameter and 1991-2020 Average (in parentheses) | Statistical/Dynamical Hybrid Forecast | Final Forecast |
|--|--|-------------------|
| Named Storms (14.4) | 16.3 | 18 |
| Named Storm Days (69.4) | 82.6 | 80 |
| Hurricanes (7.2) | 8.5 | 8 |
| Hurricane Days (27.0) | 34.2 | 35 |
| Major Hurricanes (3.2) | 4.0 | 4 |
| Major Hurricane Days (7.4) | 10.0 | 9 |
| Accumulated Cyclone Energy Index (123) | 154 | 150 |
| Net Tropical Cyclone Activity (135%) | 166 | 160 |

In addition to forecasts from ECMWF SEAS5, we continue to also incorporate a similar forecast from the UK Met Office’s GloSea6 model this year. The GloSea6 model shows comparable levels of skill to ECMWF SEAS5 at predicting the large-scale fields going into the early August statistical forecast model based on GloSea6 hindcast data from 1993-2016. Figure 10 displays observed versus hindcast ACE using the same three July predictors as used for the SEAS5 statistical/dynamical forecast model.




Figure 10: Observed versus May cross-validated statistical/dynamical hindcast values of ACE for 1993-2016 from GloSea6.

The output from the GloSea6 model also calls for a very active Atlantic hurricane season in 2021. Table 6 displays the forecasts of the three individual parameters comprising the early August statistical/dynamical hybrid forecast, while Table 7 displays the final forecast from the GloSea6 model.

Table 6: Listing of predictions of July large-scale conditions from the Met Office’s GloSea6 model output, initialized on 1 May. A plus (+) means that positive deviations of the parameter are associated with increased hurricane activity, while a minus (-) means that negative deviations of the parameter are associated with increased hurricane activity.

| Predictor | Values for 2021 Forecast | Effect on 2021 Hurricane Season |
|---|--------------------------|---------------------------------|
| 1) GloSea6 Prediction of July Surface U (10-20°N, 90-40°W) (+) | +0.3 SD | Enhance |
| 2) GloSea6 Prediction of July Skin Temperature (20-40°N, 35-15°W) (+) | +2.1 SD | Enhance |
| 3) GloSea6 Prediction of July 200 hPa U (5-15°N, 0-40°E) (-) | 0.0 SD | Neutral |

Table 7: Statistical/dynamical model output from GloSea6 for the 2021 Atlantic hurricane season and the final adjusted forecast.

| Forecast Parameter and 1991-2020 Average (in parentheses) | Statistical/Dynamical Hybrid Forecast | Final Forecast |
|--|--|-------------------|
| Named Storms (14.4) | 17.3 | 18 |
| Named Storm Days (69.4) | 89.7 | 80 |
| Hurricanes (7.2) | 9.3 | 8 |
| Hurricane Days (27.0) | 38.1 | 35 |
| Major Hurricanes (3.2) | 4.5 | 4 |
| Major Hurricane Days (7.4) | 11.4 | 9 |
| Accumulated Cyclone Energy Index (123) | 171 | 150 |
| Net Tropical Cyclone Activity (135%) | 183 | 160 |

2.3 June Analog Forecast Scheme

Certain years in the historical record have global oceanic and atmospheric trends which are similar to 2021. These years also provide useful clues as to likely levels of activity that the forthcoming 2021 hurricane season may bring. For this early June extended range forecast, we determine which of the prior years in our database have distinct trends in key environmental conditions which are similar to current May 2021 conditions and, more importantly, projected August-October 2021 conditions. Table 8 lists our analog selections.

We searched for years that were generally characterized by neutral ENSO conditions and warmer than normal North Atlantic SSTs. We anticipate that the 2021 hurricane season will have activity near the average of our six analog years.

Table 8: Analog years for 2021 with the associated hurricane activity listed for each year.

| Year | NS | NSD | H | HD | MH | MHD | ACE | NTC |
|----------------------|-----------|-----------|----------|-----------|----------|----------|------------|------------|
| 1996 | 13 | 79.00 | 9 | 45.00 | 6 | 13.00 | 166 | 192 |
| 2001 | 15 | 68.75 | 9 | 25.50 | 4 | 4.25 | 110 | 135 |
| 2006 | 10 | 58.00 | 5 | 21.25 | 2 | 2.00 | 83 | 87 |
| 2008 | 16 | 88.25 | 8 | 30.50 | 5 | 7.50 | 146 | 162 |
| 2011 | 19 | 89.75 | 7 | 26.00 | 4 | 4.50 | 126 | 145 |
| 2017 | 17 | 93.00 | 10 | 51.75 | 6 | 19.25 | 225 | 232 |
| Average | 15.0 | 79.5 | 8.0 | 33.3 | 4.5 | 8.4 | 143 | 159 |
| 2021 Forecast | 18 | 80 | 8 | 35 | 4 | 9 | 150 | 160 |

2.4 June Forecast Summary and Final Adjusted Forecast

Table 9 shows our final adjusted early June forecast for the 2021 season which is a combination of our statistical scheme, our two statistical/dynamical schemes, our

analog scheme and qualitative adjustments for other factors not explicitly contained in any of these schemes. All four of our schemes call for above-average Atlantic hurricane activity this year. Our forecast is near the average of the four schemes and calls for an above-normal season, due to both anticipated ENSO neutral conditions as well as anticipated anomalously warm SSTs in the tropical Atlantic for the peak of the Atlantic hurricane season (August-October).

Table 9: Summary of our early June statistical forecast, our two statistical/dynamical forecasts, our analog forecast, the average of those four schemes and our adjusted final forecast for the 2021 hurricane season.

| Forecast Parameter and 1991-2020 Average (in parentheses) | Statistical Scheme | SEAS5 Scheme | GloSea6 Scheme | Analog Scheme | 4-Scheme Average | Adjusted Final Forecast |
|--|-----------------------|-----------------|-------------------|------------------|---------------------|----------------------------|
| Named Storms (14.4) | 16.6 | 16.3 | 17.3 | 15.0 | 16.3 | 18 |
| Named Storm Days (69.4) | 84.2 | 82.6 | 89.7 | 79.5 | 84.0 | 80 |
| Hurricanes (7.2) | 8.7 | 8.5 | 9.3 | 8.0 | 8.6 | 8 |
| Hurricane Days (27.0) | 35.1 | 34.2 | 38.1 | 33.3 | 35.2 | 35 |
| Major Hurricanes (3.2) | 4.1 | 4.0 | 4.5 | 4.5 | 4.3 | 4 |
| Major Hurricane Days (7.4) | 10.3 | 10.0 | 11.4 | 8.4 | 10.0 | 9 |
| Accumulated Cyclone Energy Index (123) | 158 | 154 | 171 | 143 | 157 | 150 |
| Net Tropical Cyclone Activity (135%) | 170 | 166 | 183 | 159 | 170 | 160 |

3 Forecast Uncertainty

One of the questions that we are asked regarding our seasonal hurricane predictions is the degree of uncertainty that is involved. This season we continue to use probability of exceedance curves as discussed in Saunders et al. (2020). In that paper, we outlined an approach that uses statistical modeling and historical skill of various forecast models to arrive at a probability that particular values for hurricane numbers and ACE would be exceeded. Here we display probability of exceedance curves for hurricanes and ACE (Figures 11 and 12), using the error distributions calculated from both normalized cross-validated statistical as well as the cross-validated statistical/dynamical hindcasts from SEAS5. Hurricane numbers are fit to a Poisson distribution, while ACE is fit to a Weibull distribution. Table 10 displays one standard deviation uncertainty ranges (~68% of all forecasts within this range). This uncertainty estimate is also very similar to the 70% uncertainty range that NOAA provides with its forecasts. We use Poisson distributions for all storm parameters (e.g., named storms, hurricanes and major hurricanes) while we use a Weibull distribution for all integrated parameters (e.g., named storm days, ACE, etc.) except for major hurricane days We use a Laplace distribution for major hurricane days.




Figure 11: Probability of exceedance plot for hurricane numbers for the 2021 Atlantic hurricane season. The values on the x-axis indicate that the number of hurricanes exceeds that specific number. For example, 97% of Atlantic hurricane seasons from 1950-2020 have had more than two hurricanes.




Figure 12: As in Figure 11 but for ACE.

Table 10: Forecast ranges for each parameter. Note that the forecast spread may not be symmetric around the mean value, given the historical distribution of tropical cyclone activity.

| Parameter | 2021 Forecast | Uncertainty Range (68% of Forecasts Likely to Fall in This Range) |
|-------------------------------------|---------------|---|
| Named Storms (NS) | 18 | 15 – 21 |
| Named Storm Days (NSD) | 80 | 59 – 102 |
| Hurricanes (H) | 8 | 6 – 10 |
| Hurricane Days (HD) | 35 | 23 – 49 |
| Major Hurricanes (MH) | 4 | 2 – 6 |
| Major Hurricane Days (MHD) | 9 | 6 – 14 |
| Accumulated Cyclone Energy (ACE) | 150 | 100 – 205 |
| Net Tropical Cyclone (NTC) Activity | 160 | 111 – 213 |

4 ENSO

The tropical Pacific was characterized by La Niña conditions this past winter extending into early this spring, but over the past few months, the eastern and central tropical Pacific has anomalously warmed (Figure 13). NOAA officially declared La Niña over several weeks ago.

ENSO events are partially defined by NOAA based on SST anomalies in the Nino 3.4 region, which is defined as 5°S-5°N, 170-120°W. La Niña conditions are typically declared when Nino 3.4 anomalies persist at or below -0.5°C, and El Niño conditions occur when Nino 3.4 anomalies persist at or above 0.5°C. Neutral ENSO conditions span Nino 3.4 anomalies between -0.5°C and 0.5°C. Over the past two months, SST anomalies have trended upward, heralding the end of La Niña conditions and the development of neutral ENSO conditions.




Figure 13: Nino 3.4 SST anomalies from June 2020 through May 2021. Figure courtesy of Climate Prediction Center.

Upper-ocean heat content anomalies in the eastern and central tropical Pacific were at well below-normal levels from November 2020 through mid-February 2021, and since that time, upper-ocean heat content anomalies have generally increased and are now warmer than normal (Figure 14). This is another indication that the tropical Pacific has moved away from La Niña conditions.




Figure 14: Central and eastern equatorial Pacific upper ocean (0-300 meters) heat content anomalies over the past year. Upper ocean heat content anomalies have increased considerably since mid-February.

Most of the eastern and central tropical Pacific currently has SST anomalies that are near their long-term averages (Figure 15).

NOAA Coral Reef Watch Daily 5km SST Anomalies (v3.1) 29 May 2021




Figure 15: Current SST anomalies across the tropical and subtropical Pacific.

Table 11 displays March and May SST anomalies for several Nino regions. While the far eastern tropical Pacific has anomalously cooled, the rest of the eastern and central Pacific has anomalously warmed since March.

Table 11: March and May SST anomalies for Nino 1+2, Nino 3, Nino 3.4, and Nino 4, respectively. May minus March SST anomaly differences are also provided.

| Region | March SST Anomaly (°C) | May SST Anomaly (°C) | May – March SST Anomaly (°C) |
|----------|------------------------|----------------------|------------------------------|
| Nino 1+2 | -0.3 | -0.6 | -0.3 |
| Nino 3 | -0.4 | -0.4 | 0.0 |
| Nino 3.4 | -0.5 | -0.3 | +0.2 |
| Nino 4 | -0.6 | -0.1 | +0.5 |

The tropical Pacific has experienced two downwelling (warming) Kelvin waves since March which have likely helped fuel the return to ENSO neutral conditions in the tropical Pacific (Figure 16). This anomalous warming was aided by low-level westerly wind bursts.





Figure 16: Upper-ocean heat content anomalies in the tropical Pacific since June 2020. Dashed lines indicate downwelling Kelvin waves, while dotted lines indicate upwelling Kelvin waves. Downwelling Kelvin waves result in upper-ocean heat content increases, while upwelling Kelvin waves result in upper-ocean heat content decreases.

Over the next several months, we will be closely monitoring low-level winds over the tropical Pacific. Anomalous easterlies are currently observed across the central tropical Pacific, but the Climate Forecast System (CFS) is forecasting anomalous westerlies just west of the International Date Line that may promote some additional anomalous warming in the short term (Figure 17). At this point, it seems unlikely that any additional anomalous warming will be significant enough to trigger an El Niño event.



ncics.org/mjo

Mon 2021-05-31 1010 UTC

Carl Schreck
carl_schreck@ncsu.edu

Figure 17: Observed low-level winds across the equatorial region as well as predictions for the next four weeks by the Climate Forecast System. Figure courtesy of Carl Schreck.

There remains considerable uncertainty with the future state of ENSO for the peak of the Atlantic hurricane season. The latest plume of ENSO predictions from several statistical and dynamical models shows a spread for August-October, but most models are calling for neutral ENSO conditions for the next several months.




Figure 18: ENSO forecasts from various statistical and dynamical models for the Nino 3.4 SST anomaly based on late April to early May initial conditions. The majority of models are calling for ENSO neutral conditions for August-October (highlighted by the black arrow). Figure courtesy of the International Research Institute (IRI).

The latest official forecast from NOAA indicates that the chances of El Niño are quite low for August-October. NOAA is currently predicting an 8% chance of El Niño, a 50% chance of ENSO neutral conditions and a 42% chance of La Niña for the peak of the Atlantic hurricane season (Figure 19).




Figure 19: Official NOAA forecast for ENSO. The black arrow highlights the peak of the Atlantic hurricane season (August-October).

Based on the above information, our best estimate is that we will likely have neutral ENSO conditions for the peak of the Atlantic hurricane season.

5 Current Atlantic Basin Conditions

The current SST pattern across the North Atlantic basin is characterized by warm anomalies in most of the North Atlantic north of 20°N including the Gulf of Mexico, while portions of the central and eastern tropical Atlantic have near to slightly below-average SSTs (Figure 20). As we will show later, warmth in the subtropical northeastern Atlantic is typically associated with above-average Atlantic hurricane seasons.




Figure 20: Recent four-week average of North Atlantic SSTs.

The Atlantic had a relatively similar SST pattern in late March, although there are also some differences (Figure 21). Over the past two months, the subtropical Atlantic has anomalously warmed, while there has been some slight anomalous cooling in the tropical North Atlantic. Figure 22 displays SST differences between late May and late March. The North Atlantic Oscillation has predominately been in its negative phase since early April, and we have generally seen weaker trade winds, especially north of 20°N, helping to contribute to anomalous warming between 20-30°N (Figure 23). Overall, the current SST anomaly pattern correlates relatively well with what is typically seen in active Atlantic hurricane seasons (Figure 24). Anomalous warmth in the eastern Atlantic in May is typically associated with more active seasons.




Figure 21: Four-week-averaged North Atlantic SST anomalies ending on 31 March.




Figure 22: Late May 2021 minus late March 2021 SST anomaly changes.




Figure 23: Observed standardized values of the daily NAO since February. The NAO was generally positive in February and March but has been predominately negative since that time.




Figure 24: Rank correlation between May North Atlantic SST anomalies and seasonal Atlantic ACE from 1982-2020.

6 Tropical Cyclone Impact Probabilities for 2021

This year, we are debuting a new methodology for calculating the impacts of tropical cyclones for each state and county/parish along the Gulf and East Coasts, tropical cyclone-prone provinces of Canada, islands in the Caribbean and countries in Central America. We have used NOAA's Historical Hurricane Tracks [website](#) and selected all named storms, hurricanes and major hurricanes that have tracked within 50 miles of each landmass from 1851-2019. This approach allows for tropical cyclones that may have made landfall in an immediately adjacent region to be counted for all regions that were in close proximity to the landfall location of the storm. We then fit the observed frequency of storms within 50 miles of each landmass using a Poisson distribution to calculate the climatological odds of one or more events within 50 miles.

Net landfall probability is shown to be linked to the overall Atlantic basin Net Tropical Cyclone activity (NTC; see Table 12). NTC is a combined measure of the year-to-year mean of six indices of hurricane activity, each expressed as a percentage difference from the 1950-2000 climatological average. Long-term statistics show that, on average, the more active the overall Atlantic basin hurricane season is, the greater the probability of U.S. hurricane landfall.

Table 12: NTC activity in any year consists of the seasonal total of the following six parameters expressed in terms of their long-term averages. A season with 10 NS, 50 NSD, 6 H, 25 HD, 3 MH, and 5 MHD would then be the sum of the following ratios: $10/9.6 = 104$, $50/49.1 = 102$, $6/5.9 = 102$, $25/24.5 = 102$, $3/2.3 = 130$, $5/5.0 = 100$, divided by six, yielding an NTC of 107.

| 1950-2000 Average | |
|-------------------------------|------|
| 1) Named Storms (NS) | 9.6 |
| 2) Named Storm Days (NSD) | 49.1 |
| 3) Hurricanes (H) | 5.9 |
| 4) Hurricane Days (HD) | 24.5 |
| 5) Major Hurricanes (MH) | 2.3 |
| 6) Major Hurricane Days (MHD) | 5.0 |

Table 13 displays the climatological odds of storms tracking within 50 miles of each state along the Gulf and East Coasts along with the odds in 2021. Given that the seasonal forecast is for above-average hurricane activity, the odds of tropical cyclone impacts are also elevated. Probabilities for other Atlantic basin landmasses are available on our [website](#).

Table 13: Probability of ≥ 1 named storm, hurricane and major hurricane tracking within 50 miles of each coastal state from Texas to Maine. Probabilities are provided for both the 1851–2019 climatological average as well as the probability for 2021, based on the latest CSU seasonal hurricane forecast.

| State | 2021 Probability | | | Climatology | | | | |
|----------------|--|-------------|-----------|-----------------|--|-------------|-----------|-----------------|
| | Probability ≥ 1 event within 50 miles | Named Storm | Hurricane | Major Hurricane | Probability ≥ 1 event within 50 miles | Named Storm | Hurricane | Major Hurricane |
| Texas | 75% | 49% | 21% | 58% | 35% | 14% | | |
| Louisiana | 80% | 53% | 23% | 63% | 37% | 15% | | |
| Mississippi | 69% | 39% | 12% | 52% | 26% | 8% | | |
| Alabama | 75% | 41% | 14% | 58% | 28% | 9% | | |
| Florida | 96% | 75% | 41% | 86% | 58% | 28% | | |
| Georgia | 79% | 45% | 11% | 63% | 31% | 7% | | |
| South Carolina | 73% | 41% | 12% | 56% | 28% | 7% | | |
| North Carolina | 84% | 52% | 11% | 68% | 37% | 7% | | |
| Virginia | 63% | 28% | 3% | 46% | 19% | 2% | | |
| Maryland | 47% | 16% | 2% | 33% | 11% | 1% | | |
| Delaware | 33% | 8% | <1% | 22% | 5% | <1% | | |
| New Jersey | 34% | 11% | 1% | 23% | 7% | 1% | | |
| New York | 40% | 15% | 4% | 27% | 10% | 2% | | |
| Connecticut | 33% | 13% | 3% | 22% | 8% | 2% | | |
| Rhode Island | 33% | 13% | 4% | 22% | 8% | 2% | | |
| Massachusetts | 49% | 23% | 6% | 34% | 15% | 3% | | |
| New Hampshire | 28% | 10% | 3% | 18% | 6% | 2% | | |
| Maine | 34% | 12% | 3% | 23% | 8% | 2% | | |

7 Summary

An analysis of a variety of different atmosphere and ocean measurements (through May) which are known to have long-period statistical relationships with the upcoming season's Atlantic tropical cyclone activity, as well as output from dynamical models, indicate that 2021 should have above-normal activity. The big question marks with this season's predictions continue to be exactly what ENSO conditions will look like, as well as what the configuration of SSTs will look like in the Atlantic Ocean during the peak of the Atlantic hurricane season.

8 Forthcoming Updated Forecasts of 2021 Hurricane Activity

We will be issuing seasonal updates of our 2021 Atlantic basin hurricane forecasts on **Thursday 8 July, and Thursday 5 August**. We will also be issuing two-week forecasts for Atlantic TC activity during the climatological peak of the season from August–October. A verification and discussion of all 2021 forecasts will be issued in late November 2021. All of these forecasts will be available on our [website](#).

9 Verification of Previous Forecasts

Table 14: Summary verification of the authors' five previous years of seasonal forecasts for Atlantic TC activity from 2016-2020.

| 2016 | 18 April | Update 1 June | Update 1 July | Update 4 August | Obs. |
|-------------------------------|----------|------------------|------------------|--------------------|-------|
| Hurricanes | 6 | 6 | 6 | 6 | 7 |
| Named Storms | 13 | 14 | 15 | 15 | 15 |
| Hurricane Days | 21 | 21 | 21 | 22 | 27.75 |
| Named Storm Days | 52 | 53 | 55 | 55 | 81.00 |
| Major Hurricanes | 2 | 2 | 2 | 2 | 4 |
| Major Hurricane Days | 4 | 4 | 4 | 5 | 10.25 |
| Accumulated Cyclone Energy | 93 | 94 | 95 | 100 | 141 |
| Net Tropical Cyclone Activity | 101 | 103 | 105 | 110 | 155 |
| 2017 | 6 April | Update 1 June | Update 5 July | Update 4 August | Obs. |
| Hurricanes | 4 | 6 | 8 | 8 | 10 |
| Named Storms | 11 | 14 | 15 | 16 | 17 |
| Hurricane Days | 16 | 25 | 35 | 35 | 51.25 |
| Named Storm Days | 50 | 60 | 70 | 70 | 91.25 |
| Major Hurricanes | 2 | 2 | 3 | 3 | 6 |
| Major Hurricane Days | 4 | 5 | 7 | 7 | 19.25 |
| Accumulated Cyclone Energy | 75 | 100 | 135 | 135 | 226 |
| Net Tropical Cyclone Activity | 85 | 110 | 140 | 140 | 231 |
| 2018 | 5 April | Update 31 May | Update 2 July | Update 2 August | Obs. |
| Hurricanes | 7 | 6 | 4 | 5 | 8 |
| Named Storms | 14 | 14 | 11 | 12 | 15 |
| Hurricane Days | 30 | 20 | 15 | 15 | 26.75 |
| Named Storm Days | 70 | 55 | 45 | 53 | 87.25 |
| Major Hurricanes | 3 | 2 | 1 | 1 | 2 |
| Major Hurricane Days | 7 | 4 | 2 | 2 | 5.00 |
| Accumulated Cyclone Energy | 130 | 90 | 60 | 64 | 129 |
| Net Tropical Cyclone Activity | 135 | 100 | 70 | 78 | 128 |
| 2019 | 4 April | Update 4 June | Update 9 July | Update 5 August | Obs. |
| Hurricanes | 5 | 6 | 6 | 7 | 6 |
| Named Storms | 13 | 14 | 14 | 14 | 18 |
| Hurricane Days | 16 | 20 | 20 | 20 | 23.50 |
| Named Storm Days | 50 | 55 | 55 | 55 | 70.00 |
| Major Hurricanes | 2 | 2 | 2 | 2 | 3 |
| Major Hurricane Days | 4 | 5 | 5 | 5 | 9.50 |
| Accumulated Cyclone Energy | 80 | 100 | 100 | 105 | 132 |
| Net Tropical Cyclone Activity | 90 | 105 | 105 | 110 | 141 |
| 2020 | 2 April | Update 4 June | Update 7 July | Update 5 August | Obs. |
| Hurricanes | 8 | 9 | 9 | 12 | 14 |
| Named Storms | 16 | 19 | 20 | 24 | 30 |
| Hurricane Days | 35 | 40 | 40 | 45 | 34.75 |
| Named Storm Days | 80 | 85 | 85 | 100 | 118 |
| Major Hurricanes | 4 | 4 | 4 | 5 | 7 |
| Major Hurricane Days | 9 | 9 | 9 | 11 | 8.75 |
| Accumulated Cyclone Energy | 150 | 160 | 160 | 200 | 180 |
| Net Tropical Cyclone Activity | 160 | 170 | 170 | 215 | 225 |