

EXTENDED RANGE FORECAST OF ATLANTIC SEASONAL HURRICANE ACTIVITY AND LANDFALL STRIKE PROBABILITY FOR 2013

We anticipate that the 2013 Atlantic basin hurricane season will have enhanced activity compared with the 1981-2010 climatology. The tropical Atlantic has anomalously warmed over the past several months, and it appears that the chances of an El Niño event this summer and fall are unlikely. We anticipate an above-average probability for major hurricanes making landfall along the United States coastline and in the Caribbean. Coastal residents are reminded that it only takes one hurricane making landfall to make it an active season for them, and they need to prepare the same for every season, regardless of how much or how little activity is predicted.

(as of 10 April 2013)

By Philip J. Klotzbach¹ and William M. Gray²

This forecast as well as past forecasts and verifications are available via the World Wide Web at <http://hurricane.atmos.colostate.edu/Forecasts>

Emily Wilmsen, Colorado State University Media Representative, (970-491-6432) is available to answer various questions about this forecast

Department of Atmospheric Science
Colorado State University
Fort Collins, CO 80523
Email: amie@atmos.colostate.edu

¹ Research Scientist

² Professor Emeritus of Atmospheric Science

ATLANTIC BASIN SEASONAL HURRICANE FORECAST FOR 2013

Forecast Parameter and 1981-2010 Median (in parentheses)	Issue Date 10 April 2013
Named Storms (NS) (12.0)	18
Named Storm Days (NSD) (60.1)	95
Hurricanes (H) (6.5)	9
Hurricane Days (HD) (21.3)	40
Major Hurricanes (MH) (2.0)	4
Major Hurricane Days (MHD) (3.9)	9
Accumulated Cyclone Energy (ACE) (92)	165
Net Tropical Cyclone Activity (NTC) (103%)	175

PROBABILITIES FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE
LANDFALL ON EACH OF THE FOLLOWING COASTAL AREAS:

- 1) Entire U.S. coastline - 72% (average for last century is 52%)
- 2) U.S. East Coast Including Peninsula Florida - 48% (average for last century is 31%)
- 3) Gulf Coast from the Florida Panhandle westward to Brownsville - 47% (average for last century is 30%)

PROBABILITY FOR AT LEAST ONE MAJOR (CATEGORY 3-4-5) HURRICANE
TRACKING INTO THE CARIBBEAN (10-20°N, 60-88°W)

- 1) 61% (average for last century is 42%)

ABSTRACT

Information obtained through March 2013 indicates that the 2013 Atlantic hurricane season will have more activity than the median 1981-2010 season. We estimate that 2013 will have about 9 hurricanes (median is 6.5), 18 named storms (median is 12.0), 95 named storm days (median is 60.1), 40 hurricane days (median is 21.3), 4 major (Category 3-4-5) hurricanes (median is 2.0) and 9 major hurricane days (median is 3.9). The probability of U.S. major hurricane landfall is estimated to be about 140 percent of the long-period average. We expect Atlantic basin Net Tropical Cyclone (NTC) activity in 2013 to be approximately 175 percent of the long-term average

This forecast is based on a new extended-range early April statistical prediction scheme that was developed utilizing 29 years of past data. Analog predictors are also utilized. We anticipate an above-average Atlantic basin hurricane season due to the combination of an anomalously warm tropical Atlantic and a relatively low likelihood of El Niño. Coastal residents are reminded that it only takes one hurricane making landfall to make it an active season for them, and they need to prepare the same for every season, regardless of how much activity is predicted.

Why issue extended-range forecasts for seasonal hurricane activity?

We are frequently asked this question. Our answer is that it is possible to say something about the probability of the coming year's hurricane activity which is superior to climatology. The Atlantic basin has the largest year-to-year variability of any of the global tropical cyclone basins. People are curious to know how active the upcoming season is likely to be, particularly if you can show hindcast skill improvement over climatology for many past years.

Everyone should realize that it is impossible to precisely predict this season's hurricane activity in early April. There is, however, much curiosity as to how global ocean and atmosphere features are presently arranged as regards to the probability of an active or inactive hurricane season for the coming year. Our new early April statistical forecast methodology shows strong evidence over 29 past years that significant improvement over climatology can be attained. We would never issue a seasonal hurricane forecast unless we had a statistical model developed over a long hindcast period which showed significant skill over climatology.

We issue these forecasts to satisfy the curiosity of the general public and to bring attention to the hurricane problem. There is a general interest in knowing what the odds are for an active or inactive season. One must remember that our forecasts are based on the premise that those global oceanic and atmospheric conditions which preceded comparatively active or inactive hurricane seasons in the past provide meaningful information about similar trends in future seasons. This is not always true for individual seasons. It is also important that the reader appreciate that these seasonal forecasts are based on statistical schemes which, owing to their intrinsically probabilistic nature, will fail in some years. Moreover, these forecasts do not specifically predict where within the Atlantic basin these storms will strike. The probability of landfall for any one location along the coast is very low and reflects the fact that, in any one season, most U.S. coastal areas will not feel the effects of a hurricane no matter how active the individual season is.

Acknowledgment

This year's forecasts are funded by private and personal funds. We thank the GeoGraphics Laboratory at Bridgewater State University (MA) for their assistance in developing the United States Landfalling Hurricane Probability Webpage (available online at <http://www.e-transit.org/hurricane>).

The second author gratefully acknowledges the valuable input to his CSU seasonal forecast research project over many years by former graduate students and now colleagues Chris Landsea, John Knaff and Eric Blake. We also thank Professors Paul Mielke and Ken Berry of Colorado State University for statistical analysis and guidance over many years. We thank Bill Thorson for technical advice and assistance.

DEFINITIONS AND ACRONYMS

Accumulated Cyclone Energy (ACE) - A measure of a named storm's potential for wind and storm surge destruction defined as the sum of the square of a named storm's maximum wind speed (in 10^4 knots²) for each 6-hour period of its existence. The 1950-2000 average value of this parameter is 96 for the Atlantic basin.

Atlantic Multi-Decadal Oscillation (AMO) - A mode of natural variability that occurs in the North Atlantic Ocean and evidencing itself in fluctuations in sea surface temperature and sea level pressure fields. The AMO is likely related to fluctuations in the strength of the oceanic thermohaline circulation. Although several definitions of the AMO are currently used in the literature, we define the AMO based on North Atlantic sea surface temperatures from 50-60°N, 10-50°W.

Atlantic Basin - The area including the entire North Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico.

El Niño - A 12-18 month period during which anomalously warm sea surface temperatures occur in the eastern half of the equatorial Pacific. Moderate or strong El Niño events occur irregularly, about once every 3-7 years on average.

Hurricane (H) - A tropical cyclone with sustained low-level winds of 74 miles per hour (33 ms^{-1} or 64 knots) or greater.

Hurricane Day (HD) - A measure of hurricane activity, one unit of which occurs as four 6-hour periods during which a tropical cyclone is observed or is estimated to have hurricane-force winds.

Indian Ocean Dipole (IOD) - An irregular oscillation of sea surface temperatures between the western and eastern tropical Indian Ocean. A positive phase of the IOD occurs when the western Indian Ocean is anomalously warm compared with the eastern Indian Ocean.

Madden Julian Oscillation (MJO) - A globally propagating mode of tropical atmospheric intra-seasonal variability. The wave tends to propagate eastward at approximately 5 ms^{-1} , circling the globe in roughly 40-50 days.

Main Development Region (MDR) - An area in the tropical Atlantic where a majority of major hurricanes form, which we define as 7.5-22.5°N, 20-75°W.

Major Hurricane (MH) - A hurricane which reaches a sustained low-level wind of at least 111 mph (96 knots or 50 ms^{-1}) at some point in its lifetime. This constitutes a category 3 or higher on the Saffir/Simpson scale.

Major Hurricane Day (MHD) - Four 6-hour periods during which a hurricane has an intensity of Saffir/Simpson category 3 or higher.

Multivariate ENSO Index (MEI) - An index defining ENSO that takes into account tropical Pacific sea surface temperatures, sea level pressures, zonal and meridional winds and cloudiness.

Named Storm (NS) - A hurricane, a tropical storm or a sub-tropical storm.

Named Storm Day (NSD) - As in HD but for four 6-hour periods during which a tropical or sub-tropical cyclone is observed (or is estimated) to have attained tropical storm-force winds.

Net Tropical Cyclone (NTC) Activity - Average seasonal percentage mean of NS, NSD, H, HD, MH, MHD. Gives overall indication of Atlantic basin seasonal hurricane activity. The 1950-2000 average value of this parameter is 100.

Saffir/Simpson Hurricane Wind Scale - A measurement scale ranging from 1 to 5 of hurricane wind intensity. One is a weak hurricane; whereas, five is the most intense hurricane.

Southern Oscillation Index (SOI) - A normalized measure of the surface pressure difference between Tahiti and Darwin. Low values typically indicate El Niño conditions.

Sea Surface Temperature - SST

Sea Surface Temperature Anomaly - SSTA

Thermohaline Circulation (THC) - A large-scale circulation in the Atlantic Ocean that is driven by fluctuations in salinity and temperature. When the THC is stronger than normal, the AMO tends to be in its warm (or positive) phase, and more Atlantic hurricanes typically form.

Tropical Cyclone (TC) - A large-scale circular flow occurring within the tropics and subtropics which has its strongest winds at low levels; including hurricanes, tropical storms and other weaker rotating vortices.

Tropical North Atlantic (TNA) index - A measure of sea surface temperatures in the area from 5.5-23.5°N, 15-57.5°W.

Tropical Storm (TS) - A tropical cyclone with maximum sustained winds between 39 mph (18 ms^{-1} or 34 knots) and 73 mph (32 ms^{-1} or 63 knots).

Vertical Wind Shear - The difference in horizontal wind between 200 mb (approximately 40000 feet or 12 km) and 850 mb (approximately 5000 feet or 1.6 km).

1 knot = 1.15 miles per hour = 0.515 meters per second

1 Introduction

This is the 30th year in which the CSU Tropical Meteorology Project has made forecasts of the upcoming season's Atlantic basin hurricane activity. Our research team has shown that a sizable portion of the year-to-year variability of Atlantic tropical cyclone (TC) activity can be hindcast with skill exceeding climatology. This year's April forecast is based on a statistical methodology derived from 29 years of past data. Qualitative adjustments are added to accommodate additional processes which may not be explicitly represented by our statistical analyses. These evolving forecast techniques are based on a variety of climate-related global and regional predictors previously shown to be related to the forthcoming seasonal Atlantic basin TC activity and landfall probability. We believe that seasonal forecasts must be based on methods that show significant hindcast skill in application to long periods of prior data. It is only through hindcast skill that one can demonstrate that seasonal forecast skill is possible. This is a valid methodology provided that the atmosphere continues to behave in the future as it has in the past.

The best predictors do not necessarily have the best individual correlations with hurricane activity. The best forecast parameters are those that explain the portion of the variance of seasonal hurricane activity that is not associated with the other forecast variables. It is possible for an important hurricane forecast parameter to show little direct relationship to a predictand by itself but to have an important influence when included with a set of 2-3 other predictors.

A direct correlation of a forecast parameter may not be the best measure of the importance of this predictor to the skill of a 3-4 parameter forecast model. This is the nature of the seasonal or climate forecast problem where one is dealing with a very complicated atmospheric-oceanic system that is highly non-linear. There is a maze of changing physical linkages between the many variables. These linkages can undergo unknown changes from weekly to decadal time scales. It is impossible to understand how all these processes interact with each other. No one can completely understand the full complexity of the atmosphere-ocean system. But, it is still possible to develop a reliable statistical forecast scheme which incorporates a number of the climate system's non-linear interactions. Any seasonal or climate forecast scheme should show significant hindcast skill before it is used in real-time forecasts.

2 April Forecast Methodology

2.1 New April Statistical Forecast Scheme

We have developed a new April statistical forecast model which we are using for the third time this year. This model has been built over the period from 1982-2010 to incorporate the most recent and reliable data that is available. It utilizes a total of four predictors. The new Climate Forecast System Reanalysis (CFSR) (Saha et al. 2010) has been completed from 1979-present, while the NOAA Optimum Interpolation (OI) SST

(Reynolds et al. 2002) is available from 1982-present. This new model shows significant skill in predicting levels of Net Tropical Cyclone (NTC) activity over the 1982-2010 developmental period. NTC is a combined measure of the year-to-year mean of six indices of hurricane activity, each expressed as a percentage difference from the 1950-2000 climatological average. The model correlates with NTC at 0.79 when all years are included in the model, while a drop-one cross-validation (jackknife) analysis yields a correlation with NTC of 0.68. A cross-validation approach provides a more realistic view of skill the model is expected to have in future years.

Table 1 displays cross-validated NTC hindcasts for 1982-2010 along with real-time forecast values for 2011 and 2012 using the new statistical scheme, while Figure 1 displays observations versus cross-validated NTC hindcasts. We have correctly predicted above- or below-average seasons in 22 out of 31 hindcast years (71%). Our predictions have had a smaller error than climatology in 19 out of 31 years (61%). Our average hindcast error is 41 NTC units, compared with 55 NTC units for climatology. Figure 2 displays the locations of each of our predictors, while Table 2 displays the individual linear correlations between each predictor and NTC over the 1982-2010 hindcast period. All predictors correlate significantly at the 90% level using a two-tailed Student's t-test and assuming that each year represents an individual degree of freedom. The reader will note that we are incorporating a dynamical SST forecast from the European Centre for Medium-Range Weather Forecasts (ECMWF). Hindcast data provided by Frederic Vitart indicates that the ECMWF model system 3 has significant forecast skill for SSTs across the various Nino regions for September from a 1 March forecast date. We utilize the ECMWF ensemble mean prediction for September Nino 3 SSTs. The ECMWF has recently upgraded to system 4. Hindcast data from this new model is not available yet, but it is assumed that the model has improved skill to system 3. Hindcast data from 1982-2010 show that the ECMWF forecast from system 3 from a 1 March issue date correlates with observed September Nino 3 SSTs at 0.63. Table 3 displays the 2013 observed values for each of the four predictors in the new statistical forecast scheme. Table 4 displays the statistical model output for the 2013 hurricane season.

Table 1: Observed versus early April cross-validated hindcast NTC for 1982-2010 using our new forecast scheme as well as the statistical model's real-time output for 2011 and 2012. Average errors for cross-validated hindcast NTC and climatological NTC predictions are given without respect to sign. Red bold-faced years in the “Hindcast NTC” column are years that we did not go the right way, while red bold-faced years in the “Hindcast improvement over Climatology” column are years that we did not beat climatology. The hindcast went the right way with regards to an above- or below-average season in 22 out of 31 years (71%), while hindcast improvement over climatology occurred in 19 out of 31 years (61%). The hindcast has improved upon climatology in all but six years since 1993.

Year	Observed NTC	Hindcast NTC	Observed minus Hindcast	Observed minus Climatology	Hindcast improvement over Climatology
1982	38	101	-63	-62	-1
1983	31	20	11	-69	58
1984	80	163	-82	-20	-63
1985	106	60	45	6	-40
1986	37	32	5	-63	58
1987	46	71	-25	-54	29
1988	117	134	-17	17	0
1989	130	96	34	30	-4
1990	100	91	9	0	-9
1991	58	97	-39	-42	3
1992	67	20	47	-33	-14
1993	52	60	-8	-48	40
1994	35	71	-35	-65	29
1995	222	158	64	122	58
1996	192	189	3	92	89
1997	54	91	-38	-46	9
1998	169	166	3	69	66
1999	182	121	60	82	21
2000	134	154	-21	34	13
2001	135	113	22	35	13
2002	83	136	-53	-17	-36
2003	175	139	36	75	39
2004	232	89	142	132	-11
2005	279	185	94	179	85
2006	85	139	-54	-15	-39
2007	99	135	-36	-1	-35
2008	162	201	-39	62	24
2009	69	78	-9	-31	22
2010	195	235	-40	95	55
2011	145	200	-55	45	-10
2012	131	52	79	31	-46
Average	117	119	41	55	+14

Observed vs. April Model Jackknifed NTC




Figure 1: Observed versus early April jackknifed hindcast values of NTC for 1982-2010 along with real-time forecast values for 2011 and 2012.

New April Forecast Predictors




Figure 2: Location of predictors for our early April extended-range statistical prediction for the 2013 hurricane season.

Table 2: Linear correlation between each 1 April predictor and NTC over the 1982-2010 hindcast period.

Predictor	Correlation w/ NTC
1) January-March Atlantic SST (5°S-35°N, 10-40°W) (+)	0.60
2) March SLP (20-40°N, 20-35°W) (-)	-0.49
3) February-March SLP (5-20°S, 85-120°W) (+)	0.34
4) ECMWF 1 March SST Forecast for September Nino 3 (5°S-5°N, 90-150°W) (-)	-0.40

Table 3: Listing of 1 April 2013 predictors for the 2013 hurricane season. A plus (+) means that positive values of the parameter indicate increased hurricane activity.

Predictor	2013 Forecast Value	Impact on 2013 TC Activity
1) Jan-Mar Atlantic SST (5°S-35°N, 10-40°W) (+)	+0.7 SD	Increase
2) Mar SLP (20-40°N, 20-35°W) (-)	-2.9 SD	Increase
3) Feb-Mar SLP (5-20°S, 85-120°W) (+)	+0.5 SD	Increase
4) ECMWF 1 Mar SST Forecast for Sep Nino 3 (5°S-5°N, 90-150°W) (-)	+0.3 SD	Decrease

Table 4: Statistical model output for the 2013 Atlantic hurricane season, along with the final adjusted forecast.

Forecast Parameter and 1981-2010 Median (in parentheses)	Statistical Forecast	Final Forecast
Named Storms (12.0)	15.6	18
Named Storm Days (60.1)	88.9	95
Hurricanes (6.5)	9.8	9
Hurricane Days (21.3)	45.0	40
Major Hurricanes (2.0)	5.3	4
Major Hurricane Days (3.9)	14.1	9
Accumulated Cyclone Energy Index (92)	188	165
Net Tropical Cyclone Activity (103%)	200	175

2.2 Physical Associations among Predictors Listed in Table 2

The locations and brief descriptions of the predictors for our early April statistical forecast are now discussed. It should be noted that all predictors correlate with physical features during August through October that are known to be favorable for elevated levels of hurricane activity. These factors are generally related to August-October vertical wind shear in the Atlantic Main Development Region (MDR) from 10-20°N, 20-70°W as shown in Figure 3.




Figure 3: Vertical wind profile typically associated with (a) inactive Atlantic basin hurricane seasons and (b) active Atlantic basin hurricane seasons. Note that (b) has reduced levels of vertical wind shear.

For each of these predictors, we display a four-panel figure showing linear correlations between values of each predictor and August-October values of sea surface temperature (SST), sea level pressure (SLP), 200 mb zonal wind, and 850 mb zonal wind, respectively. In general, higher values of SSTs, lower values of SLP, anomalous westerlies at 850 mb and anomalous easterlies at 200 mb are associated with active Atlantic basin hurricane seasons. SST correlations are displayed using the NOAA Optimum Interpolation (OI) SST, SLP and 850 mb zonal wind correlations are displayed using the Climate Forecast System Reanalysis (CFSR), while 200 mb zonal wind correlations are displayed using the NCEP/NCAR Reanalysis, as there are questions about the quality of the upper-level wind reanalysis in the CFSR.

Predictor 1. January-March SST in the Tropical and Subtropical Eastern Atlantic (+)

(5°S-35°N, 10-40°W)

Warmer-than-normal SSTs in the tropical and subtropical Atlantic during the January-March time period are associated with a weaker-than-normal subtropical high and reduced trade wind strength during the boreal spring (Knaff 1997). Positive SSTs in January-March are correlated with weaker trade winds and weaker upper tropospheric

westerly winds, lower-than-normal sea level pressures and above-normal SSTs in the tropical Atlantic during the following August-October period (Figure 4). All three of these August-October features are commonly associated with active Atlantic basin hurricane seasons, through reductions in vertical wind shear, increased vertical instability and increased mid-tropospheric moisture, respectively. Predictor 1 correlates quite strongly (~ 0.6) with NTC. Predictor 1 also strongly correlates ($r = 0.65$) with August-October values of the Atlantic Meridional Mode (AMM) (Kossin and Vimont 2007) over the period from 1982-2010. The AMM has been shown to impact Atlantic hurricane activity through alterations in the position and intensity of the Atlantic Inter-Tropical Convergence Zone (ITCZ). Changes in the Atlantic ITCZ bring about changes in tropical Atlantic vertical and horizontal wind shear patterns and in tropical Atlantic SST patterns.

Predictor 2. March SLP in the Subtropical Atlantic (-)

(20-40°N, 20-35°W)

Our April statistical scheme in the late 1990s used a similar predictor when evaluating the strength of the March Atlantic sub-tropical ridge (Azores High). If the pressure in this area is higher than normal, it correlates strongly with increased Atlantic trade winds. These stronger trades enhance ocean mixing and upwelling, driving cooler tropical Atlantic SSTs. These cooler SSTs are associated with higher-than-normal sea level pressures which can create a self-enhancing feedback that relates to higher pressure, stronger trades and cooler SSTs during the hurricane season (Figure 5) (Knaff 1998). All three of these factors are associated with inactive hurricane seasons.

Predictor 3. February-March SLP in the southeastern tropical Pacific (+)

(5-20°S, 85-120°W)

High pressure in the southeastern tropical Pacific during the months of February-March correlates strongly with a positive Southern Oscillation Index and strong trades blowing across the eastern tropical Pacific. Strong trade winds help prevent eastward propagating Kelvin waves from transporting warmth from the western Pacific warm pool region and triggering El Niño conditions. During the August-October period, positive values of this predictor are associated with weaker trades, lower sea level pressures, and relatively cool SST anomalies in the eastern Pacific (typical of La Niña conditions) (Figure 6). The combination of these features is typically associated with more active hurricane seasons.

Predictor 4. ECMWF 1 March SST Forecast for September Nino 3 (-)


(5°S -5°N, 90-150°W)

The ECMWF seasonal forecast system 3 has shown skill at being able to predict SST anomalies associated with ENSO several months into the future (Stockdale et al. 2011). ECMWF has recently upgraded their seasonal forecast system to system 4. ENSO has been documented in many studies to be one of the primary factors associated with interannual fluctuations in Atlantic basin and U.S. landfalling hurricane activity (Gray


1984, Goldenberg and Shapiro 1996, Bove et al. 1998, Klotzbach 2011), primarily through alterations in vertical wind shear patterns. The ensemble-averaged ENSO forecast for September values of the Nino 3 region from a 1 March forecast date correlates with observations at 0.63, which is impressive considering that this forecast goes through the springtime predictability barrier, where fluctuations in ENSO lead to greatly reduced forecast skill. When the ECMWF model predicts cool SST anomalies for September, it strongly correlates with observed cool anomalies throughout the tropical Pacific associated with La Niña conditions, as well as reduced vertical wind shear, especially across the Caribbean (Figure 7).

August-October Correlations w/ Predictor 1


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U






Figure 4: Linear correlations between January-March SST in the tropical and subtropical Atlantic (Predictor 1) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). All four of these parameter deviations in the tropical Atlantic are known to be favorable for enhanced hurricane activity.

August-October Correlations w/ Predictor 2


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U






Figure 5: Linear correlations between March SLP in the subtropical Atlantic (Predictor 2) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). The predictor's primary impact during the hurricane season appears to be with MDR-averaged SST. **The correlation scale has been reversed (sign changed) to allow for easy comparison of correlations for all four predictors.**

August-October Correlations w/ Predictor 3


(a) SST



(b) SLP



(c) 850 mb U



(d) 200 mb U






Figure 6: Linear correlations between February-March SLP in the southern tropical Pacific (Predictor 3) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). The predictor's primary impacts appear to be on sea level pressure and trade wind strength across the tropical Atlantic.

August-October Correlations w/ Predictor 4 (1982-2010)

(a) SST

(b) SLP



(c) 850 mb U

(d) 200 mb U





Figure 7: Linear correlations between a 1 March ECMWF SST forecast for September Nino 3 (Predictor 4) and August-October sea surface temperature (panel a), August-October sea level pressure (panel b), August-October 850 mb zonal wind (panel c) and August-October 200 mb zonal wind (panel d). The predictor correlates very strongly with ENSO as well as vertical shear in the Caribbean. **The correlation scale has been reversed (sign changed) to allow for easy comparison of correlations for all four predictors.**

3 Forecast Uncertainty

One of the questions that we are asked regarding our seasonal hurricane predictions is the degree of uncertainty that is involved. Our predictions are our best estimate, but there is with all forecasts an uncertainty as to how well they will verify.

Table 5 provides our early April forecast, with error bars based on one standard deviation of the 1982-2010 cross-validated hindcast error. We typically expect to see 2/3 of our forecasts verify within one standard deviation of observed values, with 95% of forecasts verifying within two standard deviations of observed values. Note the rather large uncertainty ranges at this extended lead time.

Table 5: Model hindcast error and our 2013 hurricane forecast. Uncertainty ranges are given in one standard deviation (SD) increments.

Parameter	Hindcast Error (SD)	2013 Forecast	Uncertainty Range – 1 SD (67% of Forecasts Likely in this Range)
Named Storms (NS)	3.4	18	14.6 – 21.4
Named Storm Days (NSD)	21.5	95	73.5 – 116.5
Hurricanes (H)	2.4	9	5.6 – 11.4
Hurricane Days (HD)	12.7	40	27.3 – 52.7
Major Hurricanes (MH)	1.5	4	2.5 - 5.5
Major Hurricane Days (MHD)	5.5	9	3.5 - 14.5
Accumulated Cyclone Energy (ACE)	53	165	112 - 218
Net Tropical Cyclone (NTC) Activity	50	175	125 - 225

4 Analog-Based Predictors for 2013 Hurricane Activity

Certain years in the historical record have global oceanic and atmospheric trends which are similar to 2013. These years also provide useful clues as to likely trends in activity that the forthcoming 2013 hurricane season may bring. For this early April extended range forecast, we determine which of the prior years in our database have distinct trends in key environmental conditions which are similar to current February-March 2013 conditions. Table 6 lists our analog selections.

We select prior hurricane seasons since 1900 which have similar atmospheric-oceanic conditions to those currently being experienced. We searched for years that were generally characterized by neutral ENSO conditions and a negative phase of the Pacific Decadal Oscillation (PDO) during February-March, along with years that had above-average SSTs in the tropical and North Atlantic.

There were five hurricane seasons since 1900 with characteristics most similar to what we observed in February-March 2013. None of the five analog years that we selected had a significant El Niño during the peak of the hurricane season. Four out of the five years had above-average NTC activity. We anticipate that the 2013 hurricane season will have more activity than the average of our five analog years. We primarily have predicted an increase in the number of NS and NSD from the average of the five analog years, due to a likely increase in the number of weaker storms being detected

today because of advancements in satellite and aircraft technology. We believe that this season should experience well above-average activity.

Table 6: Best analog years for 2013 with the associated hurricane activity listed for each year.

Year	NS	NSD	H	HD	MH	MHD	ACE	NTC
1915	6	48.25	5	30.50	3	13.75	127	129
1952	7	39.75	6	22.75	3	7.00	87	103
1966	11	64.00	7	41.75	3	8.75	145	140
1996	13	79.00	9	45.00	6	13.00	166	192
2004	15	93.00	9	45.50	6	22.25	227	232
Average	10.4	64.8	7.2	37.1	4.2	13.0	150.6	159.2
2013 Forecast	18	95	9	40	4	9	165	175

5 ENSO

Neutral ENSO conditions were present during the winter of 2012/2013. Upper ocean heat content (top 300 meters) anomalies dropped to slightly below-normal levels during January and early February and have since returned to near average levels in the eastern and central tropical Pacific (Figure 8).




Figure 8: Central and eastern tropical Pacific upper ocean (0-300 meters) heat content anomalies over the past year. Anomalies dropped during the early portion of the winter and have since rebounded to near average levels.

Currently, SSTs are generally within 0.5°C of the average across most of the eastern and central tropical Pacific. Table 7 displays January and March SST anomalies for several Nino regions. The eastern tropical Pacific has undergone some slight warming, while the central tropical Pacific has undergone some slight cooling over the past two months.

Table 7: January and March SST anomalies for Nino 1+2, Nino 3, Nino 3.4, and Nino 4, respectively. March-January SST anomaly differences are also provided.

Region	January SST Anomaly ($^{\circ}\text{C}$)	March SST Anomaly ($^{\circ}\text{C}$)	March – January SST Anomaly ($^{\circ}\text{C}$)
Nino 1+2	-0.5	+0.2	+0.7
Nino 3	-0.5	+0.2	+0.7
Nino 3.4	-0.4	-0.2	+0.2
Nino 4	0.0	-0.3	-0.3

There is considerable uncertainty as to what is going to happen with the current neutral ENSO. The spring months are known for their ENSO predictability barrier. This is when both statistical and dynamical models show their least amount of skill. This is likely due to the fact that from a climatological perspective, trade winds across the Pacific are weakest during the late spring and early summer, and therefore, changes in phase of ENSO are often observed to occur during the April-June period. By August-October, most dynamical and statistical models are calling for the continuation of ENSO-neutral conditions (Figure 9). We find that, in general, the European Centre for Medium-Range Weather Forecasts (ECMWF) shows the best prediction skill of the various ENSO models. The correlation skill between a 1 March forecast from the ECMWF model system 3 and the observed September Nino 3.4 anomaly is 0.71, based on hindcasts/forecasts from 1982-2010, explaining half of the variance in Nino 3.4 SST. The ECMWF has recently upgraded to system 4, which is likely to have even better skill than the previous version. The hindcast skill from ECMWF is very impressive, considering that the prediction goes through the springtime predictability barrier. The average of the various ECMWF ensemble members is calling for a September Nino 3.4 SST anomaly of approximately 0.2°C . There is a fairly widespread range in the outcomes predicted by the various ensemble members, which indicates the large degree of uncertainty in future ENSO conditions (Figure 10).

Mid-Mar 2013 Plume of Model ENSO Predictions




Figure 9: ENSO forecasts from various statistical and dynamical models. Figure courtesy of the International Research Institute (IRI). Most models call for the continuation of ENSO-neutral conditions for the next few months.




Figure 10: ECMWF ensemble model forecast for the Nino 3.4 region.

The Southern Oscillation Index (SOI) has recently been somewhat above normal. The SOI is a normalized pressure differential between Tahiti and Darwin, Australia. When the SOI is positive, it implies strong trade winds across the tropical Pacific and overall, conditions typically associated with La Niña. The current SOI values are indicative of the ENSO-neutral conditions that exist across the tropical Pacific. Figure 11 displays the 30-day moving SOI since January 2011. In general, the SOI was positive through March of 2012, and has since remained relatively close to zero, with current values the highest that have been observed since mid-February of last year.




Figure 11: 30-day moving SOI since January 2011. Note how the SOI has recently been at above-average levels, indicating that a rapid transition to El Niño is unlikely.

Based on the above information, our best estimate is that we will likely remain in neutral ENSO conditions for the peak of the Atlantic hurricane season. The buildup of the warm pool in the western tropical Pacific has been relatively weak, and trade winds across the central tropical Pacific have generally been somewhat above-normal over the past few weeks. There remains a need to closely monitor ENSO conditions over the next few months. We should be more confident about ENSO conditions for the upcoming hurricane season by the time of our next forecast on June 3.

6 Current Atlantic Basin Conditions

Significant anomalous warming has occurred across the tropical Atlantic during the past two months. SSTs in the western tropical Atlantic are at near-average values, while the eastern tropical Atlantic is now significantly above average (Figure 12). Much of this anomalous warming is due to a persistent negative phase of the NAO since mid February (Figure 13). A negative phase of the NAO is associated with a weakened Atlantic subtropical high and anomalously weak trades across the tropical Atlantic. This promotes reduced mixing as well as downwelling resulting in anomalous warming. Anomalously weak westerly winds in the mid-latitudes also promote anomalous ocean currents out of the south, which contributes to general warming SSTs throughout the

North Atlantic basin. Figure 14 displays the warming in SSTs observed in the tropical Atlantic from the latter part of March minus the early part of February. The atmospheric state across the tropical Atlantic looks quite favorable for an active season, as wind shear anomalies across the basin have generally been well below average over the past two months (Figure 15).




Figure 12: March 2013 SST anomaly pattern across the Atlantic Ocean.




Figure 13: Observed NAO since December 2012. The NAO has generally been negative since mid-February.




Figure 14: Late March 2013 minus early February 2013 anomalous SST changes across the Atlantic Ocean. Note the anomalous warming that has occurred across most of the tropical Atlantic.




Figure 15: Anomalous 200-850 mb vertical wind shear from February 1 to April 1, 2013. Note the generally weaker than normal westerly shear that has been present across the tropical Atlantic. Anomalies are calculated with respect to the 1981-2010 base period and are judged to be favorable for more 2013 Atlantic hurricane activity.

7 Adjusted 2013 Forecast

Table 8 shows our final adjusted early April forecast for the 2013 season which is a combination of our statistical scheme, our analog scheme and qualitative adjustments for other factors not explicitly contained in any of these schemes. Our analog forecast calls for an active season, while the statistical model calls for a very active season.

Table 8: Summary of our early April statistical forecast, our analog forecast and our adjusted final forecast for the 2013 hurricane season.

Forecast Parameter and 1981-2010 Median (in parentheses)	Statistical Scheme	Analog Scheme	Adjusted Final Forecast
Named Storms (12.0)	15.6	10.4	18
Named Storm Days (60.1)	88.9	64.8	95
Hurricanes (6.5)	9.8	7.2	9
Hurricane Days (21.3)	45.0	37.1	40
Major Hurricanes (2.0)	5.3	4.2	4
Major Hurricane Days (3.9)	14.1	13.0	9
Accumulated Cyclone Energy Index (92)	188	151	165
Net Tropical Cyclone Activity (103%)	200	159	175

8 Landfall Probabilities for 2013

A significant focus of our recent research involves efforts to develop forecasts of the probability of hurricane landfall along the U.S. coastline and in the Caribbean. Whereas individual hurricane landfall events cannot be accurately forecast months in advance, the total seasonal probability of landfall can be forecast with statistical skill. With the observation that landfall is a function of varying climate conditions, a probability specification has been developed through statistical analyses of all U.S. hurricane and named storm landfall events during the 20th century (1900-1999). Specific landfall probabilities can be given for all tropical cyclone intensity classes for a set of distinct U.S. coastal regions.

Net landfall probability is shown linked to the overall Atlantic basin Net Tropical Cyclone activity (NTC; see Table 9). NTC is a combined measure of the year-to-year mean of six indices of hurricane activity, each expressed as a percentage difference from the 1950-2000 climatological average. Long-term statistics show that, on average, the more active the overall Atlantic basin hurricane season is, the greater the probability of U.S. hurricane landfall.

Table 9: NTC activity in any year consists of the seasonal total of the following six parameters expressed in terms of their long-term averages. A season with 10 NS, 50 NSD, 6 H, 25 HD, 3 MH, and 5 MHD would then be the sum of the following ratios: $10/9.6 = 104$, $50/49.1 = 102$, $6/5.9 = 102$, $25/24.5 = 102$, $3/2.3 = 130$, $5/5.0 = 100$, divided by six, yielding an NTC of 107.

1950-2000 Average	
1) Named Storms (NS)	9.6
2) Named Storm Days (NSD)	49.1
3) Hurricanes (H)	5.9
4) Hurricane Days (HD)	24.5
5) Major Hurricanes (MH)	2.3
6) Major Hurricane Days (MHD)	5.0

Table 10 lists landfall probabilities for the 2013 hurricane season for different TC categories for the entire U.S. coastline, the Gulf Coast and the East Coast including the Florida peninsula. We also issue probabilities for various islands and landmasses in the Caribbean and in Central America. Note that Atlantic basin NTC activity in 2013 is expected to be well above its long-term average of 100, and therefore, landfall probabilities are well above their long-term average.

Please visit the Landfalling Probability Webpage at <http://www.e-transit.org/hurricane> for landfall probabilities for 11 U.S. coastal regions and 205 coastal and near-coastal counties from Brownsville, Texas to Eastport, Maine. The probability of each U.S. coastal state being impacted by hurricanes and major hurricanes is also included. In addition, we now include probabilities of named storms, hurricanes and major hurricanes tracking within 50 and 100 miles of various islands and landmasses in the Caribbean and Central America. We suggest that all coastal residents visit the Landfall Probability Webpage for their individual probabilities.

As an example we find that the probability of Florida being hit by a major (Cat 3-4-5) hurricane this year is 34% which is well above the climatological average of 21%.

South Florida is much more prone to being impacted by a hurricane on an individual year basis compared with northeast Florida. For instance, the probability of Miami-Dade County being impacted by hurricane-force wind gusts this year is 19%. For Duval County, the probability of being impacted by hurricane-force wind gusts is only 5%. However, considering a 50-year period, the probability of Duval County experiencing hurricane-force wind gusts is 75%.

For the island of Puerto Rico, the probability of a named storm, hurricane and major hurricane tracking within 50 miles of the island this year is 50%, 26%, and 8%, respectively.

Table 10: Estimated probability (expressed in percent) of one or more landfalling tropical storms (TS), category 1-2 hurricanes (HUR), category 3-4-5 hurricanes, total hurricanes and named storms along the entire U.S. coastline, along the Gulf Coast (Regions 1-4), and along the Florida Peninsula and the East Coast (Regions 5-11) for 2013.

Probabilities of a tropical storm, hurricane and major hurricane tracking into the Caribbean are also provided. The long-term mean annual probability of one or more landfalling systems during the last 100 years is given in parentheses.

Region	TS	Category 1-2 HUR	Category 3-4-5 HUR	All HUR	Named Storms
Entire U.S. (Regions 1-11)	94% (79%)	86% (68%)	72% (52%)	96% (84%)	99% (97%)
Gulf Coast (Regions 1-4)	79% (59%)	62% (42%)	47% (30%)	80% (60%)	96% (83%)
Florida plus East Coast (Regions 5-11)	71% (50%)	64% (44%)	48% (31%)	81% (61%)	94% (81%)
Caribbean (10-20°N, 60-88°W)	95% (82%)	77% (57%)	61% (42%)	91% (75%)	99% (96%)

9 Summary

An analysis of a variety of different atmosphere and ocean measurements (through March) which are known to have long-period statistical relationships with the upcoming season's Atlantic tropical cyclone activity indicate that 2013 should be a very active hurricane season. The only apparent obstacle to this assessment would be the formation of a moderate to strong El Niño event which we currently judge to be unlikely.

10 Can Rising Levels of CO₂ be Associated with the Devastation caused by Hurricane Sandy (2012) along with the Increase in Atlantic Hurricane Activity since 1995?

We have extensively discussed this topic in many previous papers which can be found on our Tropical Meteorology website. For more information on this topic we refer you to the following three references, which can be accessed by clicking on the links below:

[Gray, W. M., and P. J. Klotzbach, 2012: US Hurricane Damage - Can Rising Levels of CO₂ be Associated with Sandy's Massive Destruction? *Colorado State University Publication*, 23 pp.](#)

[W. M. Gray, and P. J. Klotzbach, 2013: Tropical cyclone forecasting. **National Hurricane Conference**, New Orleans, Louisiana, March 28, 2013.](#)

[W. M. Gray, and P. J. Klotzbach, 2013: Wind destruction from hurricanes. **Windstorm Insurance Conference**, Orlando, Florida, January 30, 2013.](#)

11 Forthcoming Updated Forecasts of 2013 Hurricane Activity

We will be issuing seasonal updates of our 2013 Atlantic basin hurricane forecasts on **Monday 3 June, and Friday 2 August**. We will also be issuing two-week forecasts for Atlantic TC activity during the climatological peak of the season from August-October. A verification and discussion of all 2013 forecasts will be issued in late November 2013. All of these forecasts will be available on the web at: <http://hurricane.atmos.colostate.edu/Forecasts>.

12 Acknowledgments

Besides the individuals named on page 5, there have been a number of other meteorologists that have furnished us with data and given valuable assessments of the current state of global atmospheric and oceanic conditions. These include Brian McNoldy, Art Douglas, Ray Zehr, Mark DeMaria, Todd Kimberlain, Paul Roundy and Amato Evan. In addition, Barbara Brumit and Amie Hedstrom have provided excellent manuscript, graphical and data analysis and assistance over a number of years. We have profited over the years from many in-depth discussions with most of the current and past NHC hurricane forecasters. The second author would further like to acknowledge the encouragement he has received for this type of forecasting research application from Neil Frank, Robert Sheets, Robert Burpee, Jerry Jarrell, Max Mayfield, and Bill Read former directors of the National Hurricane Center (NHC), and the current director, Rick Knabb.

13 Citations and Additional Reading

- Alexander, M. A., I. Blade, M. Newman, J. R. Lanzante, N.-C. Lau, and J. D. Scott, 2002: The atmospheric bridge: The influence of ENSO teleconnections on air-sea interaction over the global oceans. *J. Climate*, 15, 2205-2231.
- Blake, E. S., 2002: Prediction of August Atlantic basin hurricane activity. Dept. of Atmos. Sci. Paper No. 719, Colo. State Univ., Ft. Collins, CO, 80 pp.
- Blake, E. S. and W. M. Gray, 2004: Prediction of August Atlantic basin hurricane activity. *Wea. Forecasting*, 19, 1044-1060.
- Chiang, J. C. H. and D. J. Vimont, 2004: Analogous Pacific and Atlantic meridional modes of tropical atmosphere-ocean variability. *J. Climate*, 17, 4143-4158.
- DeMaria, M., J. A. Knaff and B. H. Connell, 2001: A tropical cyclone genesis parameter for the tropical Atlantic. *Wea. Forecasting*, 16, 219-233.
- Elsner, J. B., G. S. Lehmiller, and T. B. Kimberlain, 1996: Objective classification of Atlantic hurricanes. *J. Climate*, 9, 2880-2889.
- Evan, A. T., J. Dunion, J. A. Foley, A. K. Heidinger, and C. S. Velden, 2006: New evidence for a relationship between Atlantic tropical cyclone activity and African dust outbreaks, *Geophys. Res. Lett*, 33, doi:10.1029/2006GL026408.
- Goldenberg, S. B., C. W. Landsea, A. M. Mestas-Nunez, and W. M. Gray, 2001: The recent increase in Atlantic hurricane activity: Causes and Implications. *Science*, 293, 474-479.
- Goldenberg, S. B. and L. J. Shapiro, 1996: Physical mechanisms for the association of El Niño and West African rainfall with Atlantic major hurricane activity. *J. Climate*, 1169-1187.
- Gray, W. M., 1984a: Atlantic seasonal hurricane frequency: Part I: El Niño and 30 mb quasi-biennial oscillation influences. *Mon. Wea. Rev.*, 112, 1649-1668.
- Gray, W. M., 1984b: Atlantic seasonal hurricane frequency: Part II: Forecasting its variability. *Mon. Wea. Rev.*, 112, 1669-1683.
- Gray, W. M., 1990: Strong association between West African rainfall and US landfall of intense hurricanes. *Science*, 249, 1251-1256.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1992: Predicting Atlantic seasonal hurricane activity 6-11 months in advance. *Wea. Forecasting*, 7, 440-455.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1993: Predicting Atlantic basin seasonal tropical cyclone activity by 1 August. *Wea. Forecasting*, 8, 73-86.
- Gray, W. M., C. W. Landsea, P. W. Mielke, Jr., and K. J. Berry, 1994a: Predicting Atlantic basin seasonal tropical cyclone activity by 1 June. *Wea. Forecasting*, 9, 103-115.
- Gray, W. M., J. D. Sheaffer and C. W. Landsea, 1996: Climate trends associated with multi-decadal variability of intense Atlantic hurricane activity. Chapter 2 in "Hurricanes, Climatic Change and Socioeconomic Impacts: A Current Perspective", H. F. Diaz and R. S. Pulwarty, Eds., Westview Press, 49 pp.

- Gray, W. M., 1998: Atlantic ocean influences on multi-decadal variations in El Niño frequency and intensity. Ninth Conference on Interaction of the Sea and Atmosphere, 78th AMS Annual Meeting, 11-16 January, Phoenix, AZ, 5 pp.
- Grossmann, I. and P. J. Klotzbach, 2009: A review of North Atlantic modes of natural variability and their driving mechanisms. *J. Geophys. Res.*, 114, D24107, doi:10.1029/2009JD012728.
- Henderson-Sellers, A., H. Zhang, G. Berz, K. Emanuel, W. Gray, C. Landsea, G. Holland, J. Lighthill, S.-L. Shieh, P. Webster, and K. McGuffie, 1998: Tropical cyclones and global climate change: A post-IPCC assessment. *Bull. Amer. Meteor. Soc.*, 79, 19-38.
- Klotzbach, P. J., 2002: Forecasting September Atlantic basin tropical cyclone activity at zero and one-month lead times. Dept. of Atmos. Sci. Paper No. 723, Colo. State Univ., Ft. Collins, CO, 91 pp.
- Klotzbach, P. J., 2006: Trends in global tropical cyclone activity over the past twenty years (1986-2005). *Geophys. Res. Lett.*, 33, doi:10.1029/2006GL025881.
- Klotzbach, P. J., 2007: Revised prediction of seasonal Atlantic basin tropical cyclone activity from 1 August. *Wea. and Forecasting*, 22, 937-949.
- Klotzbach, P. J. and W. M. Gray, 2003: Forecasting September Atlantic basin tropical cyclone activity. *Wea. and Forecasting*, 18, 1109-1128.
- Klotzbach, P. J. and W. M. Gray, 2004: Updated 6-11 month prediction of Atlantic basin seasonal hurricane activity. *Wea. and Forecasting*, 19, 917-934.
- Klotzbach, P. J. and W. M. Gray, 2006: Causes of the unusually destructive 2004 Atlantic basin hurricane season. *Bull. Amer. Meteor. Soc.*, 87, 1325-1333.
- Knaff, J. A., 1997: Implications of summertime sea level pressure anomalies. *J. Climate*, 10, 789-804.
- Knaff, J. A., 1998: Predicting summertime Caribbean sea level pressure. *Wea. and Forecasting*, 13, 740-752.
- Kossin, J. P., and D. J. Vimont, 2007: A more general framework for understanding Atlantic hurricane variability and trends. *Bull. Amer. Meteor. Soc.*, 88, 1767-1781.
- Landsea, C. W., 1991: West African monsoonal rainfall and intense hurricane associations. Dept. of Atmos. Sci. Paper, Colo. State Univ., Ft. Collins, CO, 272 pp.
- Landsea, C. W., 1993: A climatology of intense (or major) Atlantic hurricanes. *Mon. Wea. Rev.*, 121, 1703-1713.
- Landsea, C. W., 2007: Counting Atlantic tropical cyclones back to 1900. *EOS*, 88, 197, 202.
- Landsea, C. W. and W. M. Gray, 1992: The strong association between Western Sahel monsoon rainfall and intense Atlantic hurricanes. *J. Climate*, 5, 435-453.
- Landsea, C. W., W. M. Gray, P. W. Mielke, Jr., and K. J. Berry, 1992: Long-term variations of Western Sahelian monsoon rainfall and intense U.S. landfalling hurricanes. *J. Climate*, 5, 1528-1534.
- Landsea, C. W., W. M. Gray, K. J. Berry and P. W. Mielke, Jr., 1996: June to September rainfall in the African Sahel: A seasonal forecast for 1996. 4 pp.
- Landsea, C. W., N. Nicholls, W.M. Gray, and L.A. Avila, 1996: Downward trends in the frequency of intense Atlantic hurricanes during the past five decades. *Geo. Res. Letters*, 23, 1697-1700.

- Landsea, C. W., R. A. Pielke, Jr., A. M. Mestas-Nunez, and J. A. Knaff, 1999: Atlantic basin hurricanes: Indices of climatic changes. *Climatic Changes*, 42, 89-129.
- Landsea, C.W. et al., 2005: Atlantic hurricane database re-analysis project. Available online at http://www.aoml.noaa.gov/hrd/data_sub/re_anal.html
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1996: Artificial skill and validation in meteorological forecasting. *Wea. Forecasting*, 11, 153-169.
- Mielke, P. W., K. J. Berry, C. W. Landsea and W. M. Gray, 1997: A single sample estimate of shrinkage in meteorological forecasting. *Wea. Forecasting*, 12, 847-858.
- Pielke, Jr. R. A., and C. W. Landsea, 1998: Normalized Atlantic hurricane damage, 1925-1995. *Wea. Forecasting*, 13, 621-631.
- Pielke, Jr. R. A., and J. Gratz, C. W. Landsea, D. Collins, and R. Masulin, 2008: Normalized hurricane damage in the United States: 1900-2005. *Nat. Haz. Rev.*, 9, 29-42, doi:10.1061/(ASCE)1527-6988(2008)9:1(29).
- Rasmusson, E. M. and T. H. Carpenter, 1982: Variations in tropical sea-surface temperature and surface wind fields associated with the Southern Oscillation/El Niño. *Mon. Wea. Rev.*, 110, 354-384.
- Seseske, S. A., 2004: Forecasting summer/fall El Niño-Southern Oscillation events at 6-11 month lead times. Dept. of Atmos. Sci. Paper No. 749, Colo. State Univ., Ft. Collins, CO, 104 pp.
- Vimont, D. J., and J. P. Kossin, 2007: The Atlantic meridional mode and hurricane activity. *Geophys. Res. Lett.*, 34, L07709, doi:10.1029/2007GL029683.
- Wheeler, M. C., and H. H. Hendon, 2004: An all-season real-time multivariate MJO index: Development of an index for monitoring and prediction. *Mon. Wea. Rev.*, 132, 1917-1932.

14 Verification of Previous Forecasts

Table 11: Summary verification of the authors' four previous years of seasonal forecasts for Atlantic TC activity between 2009-2012. Verifications of all seasonal forecasts back to 1984 are available here: http://tropical.atmos.colostate.edu/Includes/Documents/Publications/forecast_verifications.xls

2009	10 Dec. 2008	Update 9 April	Update 2 June	Update 4 August	Obs.
Hurricanes	7	6	5	4	3
Named Storms	14	12	11	10	9
Hurricane Days	30	25	20	18	12
Named Storm Days	70	55	50	45	30
Major Hurricanes	3	2	2	2	2
Major Hurricane Days	7	5	4	4	3.50
Accumulated Cyclone Energy	125	100	85	80	53
Net Tropical Cyclone Activity	135	105	90	85	69

2010	9 Dec. 2009	Update 7 April	Update 2 June	Update 4 August	Obs.
Hurricanes	6-8	8	10	10	12
Named Storms	11-16	15	18	18	19
Hurricane Days	24-39	35	40	40	38.50
Named Storm Days	51-75	75	90	90	89.50
Major Hurricanes	3-5	4	5	5	5
Major Hurricane Days	6-12	10	13	13	11
Accumulated Cyclone Energy	100-162	150	185	185	165
Net Tropical Cyclone Activity	108-172	160	195	195	196

2011	8 Dec. 2010	Update 6 April	Update 1 June	Update 3 August	Obs.
Hurricanes	9	9	9	9	7
Named Storms	17	16	16	16	19
Hurricane Days	40	35	35	35	26
Named Storm Days	85	80	80	80	89.75
Major Hurricanes	5	5	5	5	4
Major Hurricane Days	10	10	10	10	4.5
Net Tropical Cyclone Activity	180	175	175	175	145

2012	4 April	Update 1 June	Update 3 August	Obs.
Hurricanes	4	5	6	10
Named Storms	10	13	14	19
Hurricane Days	16	18	20	28.,50
Named Storm Days	40	50	52	101
Major Hurricanes	2	2	2	2
Major Hurricane Days	3	4	5	0.50
Accumulated Cyclone Energy	70	80	99	133
Net Tropical Cyclone Activity	75	90	105	131